

Light supersymmetric axion in an anomalous Abelian extension of the standard modelClaudio Corianò,^{*} Marco Guzzi,[†] Antonio Mariano,[‡] and Simone Morelli[§]*Dipartimento di Fisica, Università del Salento Via Arnesano 73100 Lecce, Italy**and INFN Sezione di Lecce, Via Arnesano 73100 Lecce, Italy*

(Received 12 January 2009; published 10 August 2009)

We present a supersymmetric extension of the standard model (USSM-A) with an anomalous $U(1)$ and Stückelberg axions for anomaly cancellation, generalizing similar nonsupersymmetric constructions. The model, built by a bottom-up approach, is expected to capture the low-energy supersymmetric description of axionic symmetries in theories with gauged anomalous Abelian interactions, previously explored in the nonsupersymmetric case for scenarios with intersecting branes. The choice of a USSM-like superpotential, with one extra singlet superfield and an extra Abelian symmetry, allows a physical axionlike particle in the spectrum. We describe some general features of this construction and, in particular, the modification of the dark-matter sector which involves both the axion and several neutralinos with an axino component. The axion is expected to be very light in the absence of phases in the superpotential but could acquire a mass which can also be in the few GeV range or larger. In particular, the gauging of the anomalous symmetry allows independent mass/coupling interaction to the gauge fields of this particle, a feature which is absent in traditional (invisible) axion models. We comment on the general implications of our study for the signature of moduli from string theory due to the presence of these anomalous symmetries.

DOI: 10.1103/PhysRevD.80.035006

PACS numbers: 12.60.Jv

I. INTRODUCTION

Extensions of the standard model (SM) describing axionlike particles—and with supersymmetry as a basic low-energy scenario—are an interesting area of investigation which has the potentiality to provide an answer to a series of unsolved theoretical issues. Among them are those concerning the possible presence of anomalous extra neutral gauge interactions at current and future colliders in some special channels, especially in the search for an anomalous extra Z' . This investigation could also clarify the role of weakly coupled pseudoscalars in the early universe. For this reason several studies addressing the experimental detection of pseudoscalars at future experiments [1–6], has received an impressive impulse in the recent literature.

One of the distinctive features of these extensions is the presence of extra Abelian interactions which are anomalous. We just recall that anomalous $U(1)$'s are quite common in several string constructions and that the mechanism of anomaly cancellation, if realized at low-energy by a Wess-Zumino counterterm (WZ), may cause the presence of a physical axion in the spectrum. This result points directly towards the possibility of having a new dark-matter candidate (see also [7]), which is certainly one of the most appealing features of this class of theories [8].

One of the first successful realization of the nonsupersymmetric version of these models comes from special

vacua of string/brane theory (orientifold vacua), in the form of stacks of intersecting branes, which induce a gauge structure given by the product of $U(N) \sim SU(N) \times U(1)$ factors, where N is the number of branes of each stack (see [9] for an overview). Among the $U(1)$ factors, one of them is identified with the SM hypercharge ($U(1)_Y$), while the remaining ones are anomalous and involve Stückelberg axions for anomaly cancellation. In effective string models the Abelian structure is in general characterized by the presence of several $U(1)$ factors, described in the hypercharge basis by direct products of the form $G_1 \equiv U(1)_Y \times U(1)_1 \times \dots \times U(1)_p$, with an anomaly-free hypercharge generator and p anomalous $U(1)$'s which are accompanied by axions b_i , with $i = 1, 2, \dots, p$. The anomalous $U(1)$'s in this construction are in a broken phase, called the “Stückelberg phase.” In particular, after electroweak symmetry breaking (EWSB), one of the axions becomes physical [8] and is characterized by independent mass/coupling relations, where the coupling appears in an ordinary $b\bar{F}\bar{F}$ interaction with the gauge fields, providing a generalization of the Peccei-Quinn (PQ) axion. One shortcoming of this description, at this time, is the absence of a supersymmetric extension of it with the appearance of a physical axion. The generalization to the supersymmetric case of these theories is interesting on several grounds. For instance, it allows to study an entire new class of extensions of the MSSM in the presence of a gauging of the axionic symmetries [10] and, at the same time, represents an intermediate step toward the unification with gravity of the same models, within certain formulations of supergravity [11,12]. The formulation of [10], which is specific for a MSSM superpotential parallels a previous general study of the same topic contained in [11].

^{*}claudio.coriano@le.infn.it[†]marco.guzzi@le.infn.it[‡]antonio.mariano@le.infn.it[§]simone.morelli@le.infn.it

Therefore, these types of constructions provide a consistent framework for the study of the effects of moduli of string/brane theory within scenarios with large extra dimensions or via supergravity, together with their low-energy implications in cosmology and in collider physics [13]. Recently, an extension of the MSSM containing an anomalous $U(1)$, made massive by a Stückelberg supermultiplet [14] has been introduced in [10]. This has been based on the superpotential of the MSSM with an extra Abelian symmetry. One of the features of this construction is the absence of a Higgs-axion mixing, since the bosonic component of the Stückelberg multiplet remains an ordinary Goldstone mode. Therefore, the final theory is characterized by a physical axino but not by a physical axion. The objective of our analysis is to show that a similar construction can be performed in more general ways, thereby generating a model with a physical axionlike particle. This provides a complete supersymmetric generalization of the (gauged) PQ axion. We will work out the requirements that are needed in order to make this extension possible, detailing some of the arguments that have been presented in short form in [15] and analyzing the main features of the effective action of such a theory, that we call the USSM-A due to the anomalous $U(1)$ (A) and to the specific choice of the USSM superpotential.

Our work is organized as follows. We briefly describe the class of models that we are going to investigate, outlining their basic structure, together with their supersymmetric generalizations. Along the way, we will underline the differences between our construction and the previous construction of [10]. We show how a physical axion is bound to appear in the spectrum and describe all the sectors of this theory. We derive the corresponding generalized Ward identities and characterize the Chern-Simons interactions of this class of models bringing up one typical example of application. We study the neutralino sector of the model and present a brief numerical analysis of its spectrum. Most of our attention in this work focuses on the basic characterization of this model, stressing on the mechanism that allows a physical axion in the spectrum. We conclude with some comments on possible extensions of this analysis to more general potentials characterized by moduli in different scenarios derived from string theory.

II. SUPERSYMMETRIC EXTENSIONS OF THE STANDARD MODEL AND EXTRA $U(1)$ 'S

Abelian (anomaly-free) supersymmetric extensions of the SM have been discussed in several previous works [16–21]. In [20] the authors explore an extension of the minimal supersymmetric SM (MSSM) with an extra SM singlet chiral superfield \hat{S} , with chiral charges chosen so to allow trilinear couplings of \hat{S} to the two Higgs doublets \hat{H}_1 , \hat{H}_2 in the superpotential. The μ term, in this case, is generated by the vacuum expectation value (vev) of the

scalar component of \hat{S} , precisely by the $\hat{S}\hat{H}_1 \cdot \hat{H}_2$ interaction. The structure of this model, usually called USSM, shares some similarity with the nearly-minimal supersymmetric SM (nMSSM) [22] and the next-to-minimal supersymmetric SM (NMSSM) [23]. In all of these three models the extra scalar \hat{S} is introduced for the same purpose but in the nMSSM and NMSSM this field is a singlet under the complete gauge group (which is the same as the SM) while in the USSM the field is charged under the extra $U(1)$. We recall that the nMSSM and the NMSSM differ at the level of the superpotential in the structure of the pure \hat{S} contribution, which is either linear (nMSSM) or cubic (NMSSM).

In the approach of [20] this appears to be a necessary requirement since a scalar superfield, singlet under the complete $SU(3) \times SU(2) \times U(1)_Y \times U(1)_B$ gauge group, while solving the μ problem, however, does not allow a consistent pattern of EWSB, leaving the extra Z' of the neutral sector massless. This construction is realized with an anomaly-free chiral spectrum.

A. MSSM and USSM with an anomalous $U(1)$ '

In [10] the authors investigate a supersymmetric extension of the SM with an extra $U(1)$, based on the superpotential of the MSSM. They make an important step forward in the analysis of this class of theories, using a bottom-up approach, that is by (1) fixing the effective action of the anomalous Abelian symmetry using the Stückelberg supermultiplet to give mass to the anomalous gauge boson and (2) using Wess-Zumino counterterms to balance the mixed and cubic $U(1)_B$ anomalies of the theory. A third element of the construction is the possible presence of Chern-Simons interactions [8] which find their way to low energy from string theory [24], and which amount to a redistribution of the anomaly starting from a symmetric distribution on each leg of the anomaly vertex. This redistribution is allowed whenever the symmetry of the vertex does not allow to uniquely define the breaking of the Ward identities separately on each of its legs. The meaning of this freedom, from the point of view of the effective field theory, is that each model allows a set of additional (defining) Ward identities for the distribution of the anomaly which are a specific feature of anomalous models in which the trilinear gauge interactions are not identically zero (in the massless fermion phase, the chiral phase).

In the first supersymmetric version of these models [10], the ordinary MSSM Lagrangian is naturally extended by the Stückelberg multiplet which provides a kinetic term for the same multiplet while rendering the extra Z' massive. The defining phase of the model is, therefore, the Stückelberg phase. In this construction the bosonic partner of the axino, which is the fermionic component of the multiplet, remains a Goldstone mode after EWSB and is therefore unphysical.

B. Inducing Higgs-axion mixing

At the origin of the physical axion is the mechanism of Higgs-axion mixing. For this to take place one needs a Higgs sector which is charged under the anomalous $U(1)_B$ so that the mass of the anomalous gauge boson comes from a combination of the Higgs and Stückelberg mechanisms. In the case of the MSSM this mixing does not occur even if the two Higgses are charged under the anomalous $U(1)$. The presence of a μ term in the superpotential forces the two charges of the two Higgs doublets to take opposite values, thereby guaranteeing also the cancellation of the extra anomalies due to the circulating Higgsinos, but is not enough to give mass to the anomalous gauge boson. In other words, in the absence of a Stückelberg multiplet the mass matrix of the gauge boson has still an additional null eigenvalue. The true mechanism of mass generation of the anomalous Z' , therefore, is just the Stückelberg, which in this situation is a Goldstone mode. In fact, one reobtains a massive Yang-Mills theory just by going to the unitary gauge and eliminating the axion.

III. THE STRUCTURE OF THE MODEL

A simple way out in order to have Higgs-axion mixing and a light axion in the physical spectrum consists to use a modified superpotential as in [20], but now with an anomalous gauge structure, and to combine it with the Lagrangian of the Stückelberg supermultiplet. In other words, we move from the superpotential of MSSM-type to the one typical of the USSM, introducing an extra scalar superfield \hat{S} which is nonsinglet under an extra $U(1)_B$, maintaining the anomalous structure induced by the extra neutral current. This specific assumption allows to remove the second massless eigenvalue in the mass matrix of the gauge bosons and allows to induce Higgs-axion mixing once that the Stückelberg mechanism is invoked to contribute to the mass of the extra Z' . The conditions that need to be verified in order to have a physical axion in the spectrum are obtained from an analysis of the CP -odd sector of the theory and involve both the potential and the derivative couplings (mixings) of the massive gauge bosons with their Goldstones ($Z_I \partial G_{Z_I}$) extracted from the broken phase. In general, the presence of extra singlet superfields in the superpotential allows such a mixing and we will illustrate this requirement in one of the sections below. The analysis that we will present in the next sections has the goal to clarify this point, starting from the MSSM case, where none of the CP -odd states acquires an axionlike coupling.

These new features do not affect the chargino sector with respect to the MSSM.

IV. THE SUPERPOTENTIAL

The construction of models characterized by a physical axion in their spectrum requires an appropriate superpotential. In order to obtain this, we consider the introduction

of an extra SM singlet \hat{S} . For this reason, the superpotential of the model investigated is given by

$$\mathcal{W} = \lambda \hat{S} \hat{H}_1 \cdot \hat{H}_2 + y_e \hat{H}_1 \cdot \hat{L} \hat{R} + y_d \hat{H}_1 \cdot \hat{Q} \hat{D}_R + y_u \hat{H}_2 \cdot \hat{Q} \hat{U}_R, \quad (1)$$

which coincides with the model of [20], called the USSM. We refer to Table I for a list of the charge assignment of the chiral superfields of our model and Table III (see appendix) for the list of their bosonic and fermionic components; the scalar superfields corresponding to $SU(3)$, $SU(2)$, $U(1)_Y$ and $U(1)_B$ are, respectively, $\hat{G}^a(x, \theta, \bar{\theta})$ (with $a = 1, 2, \dots, 8$), $\hat{W}^i(x, \theta, \bar{\theta})$ (with $i = 1, 2, 3$), $\hat{Y}(x, \theta, \bar{\theta})$ and $\hat{B}(x, \theta, \bar{\theta})$ and they fall in the usual adjoint representations of the gauge group factors.

We have denoted the charges by $Q_{f,X}$, where X denotes the hypercharge (Y), the charged W^\pm bosons (W), the non-Abelian gluons (G) and the anomalous gauge boson (B). At the same time we denote with B_X the charges of the X superfield respect to the anomalous $U(1)$. Unlike the NMSSM and the nMSSM, \mathcal{W} does not contain linear and cubic terms in \hat{S} in order to preserve the gauge invariance in the presence of a nonvanishing B_S charge. This requirement is strictly necessary if the extra scalar \hat{S} is only a SM singlet. Gauge invariance gives the conditions

$$\begin{aligned} B_{H_1} + B_{H_2} + B_S = 0 & & B_{H_1} + B_L + B_R = 0 \\ B_{H_1} + B_Q + B_{D_R} = 0 & & B_{H_2} + B_Q + B_{U_R} = 0, \end{aligned} \quad (2)$$

which will be used below. It is not hard to show that the possibility of declaring \hat{S} to be a singlet under the entire gauge group ($B_S = 0$) $SU(3) \times SU(2) \times G_1$ leaves an extra gauge boson massless beside the photon, after EWSB and as such it is not acceptable.

Anomaly cancellation: Defining the model

We start by identifying the anomalous contributions of the model, whose gauge structure is of the form $SU(3) \times SU(2) \times U(1)_Y \times U(1)_B$.

The anomalous trilinear gauge interactions are all the ones involving the extra anomalous $U(1)_B$, namely $\{U(1)_B, U(1)_B, U(1)_B\}$, $\{U(1)_B, U(1)_Y, U(1)_Y\}$, $\{U(1)_B, U(1)_B,$

TABLE I. Charge assignment of the model; boldface numbers indicate the dimensions of the corresponding representations.

Superfields	$SU(3)$	$SU(2)$	$U(1)_Y$	$U(1)_B$
$\hat{\mathbf{b}}(x, \theta, \bar{\theta})$	1	1	0	\dots
$\hat{S}(x, \theta, \bar{\theta})$	1	1	0	B_S
$\hat{L}(x, \theta, \bar{\theta})$	1	2	-1/2	B_L
$\hat{R}(x, \theta, \bar{\theta})$	1	1	1	B_R
$\hat{Q}(x, \theta, \bar{\theta})$	3	2	1/6	B_Q
$\hat{U}_R(x, \theta, \bar{\theta})$	$\bar{3}$	1	-2/3	B_{U_R}
$\hat{D}_R(x, \theta, \bar{\theta})$	$\bar{3}$	1	+1/3	B_{D_R}
$\hat{H}_1(x, \theta, \bar{\theta})$	1	2	-1/2	B_{H_1}
$\hat{H}_2(x, \theta, \bar{\theta})$	1	2	1/2	B_{H_2}

$U(1)_Y$, $\{U(1)_B, SU(2), SU(2)\}$, $\{U(1)_B, SU(3), SU(3)\}$. In terms of the charges we can write each sector as follows

$$\begin{aligned}\mathcal{A}_{BBB} &= \sum_f Q_{f,B}^3 & \mathcal{A}_{BYY} &= \sum_f Q_{f,B} Q_{f,Y}^2 \\ \mathcal{A}_{BBY} &= \sum_f Q_{f,B}^2 Q_{f,Y} & \mathcal{A}_{BWW} &= \sum_f Q_{f,B} \text{Tr}[\tau^i \tau^j] \\ \mathcal{A}_{BGG} &= \sum_f Q_{f,B} \text{Tr}[T^a T^b],\end{aligned}\quad (3)$$

where T^a are the generators of $SU(3)$ and τ^i the Pauli matrices. Compared to the analysis of [10], here we have anomalous trilinear interactions also in the sector involving the $SU(3)$ mixed anomaly due to the nonvanishing charge B_S . Using the constraints coming from the Yukawa couplings and the conditions of gauge invariance, the expressions of the anomalies take the form

$$\begin{aligned}\mathcal{A}_{BBB} &= 3(6B_Q^3 + 3B_{U_R}^3 + 3B_{D_R}^3) + (6B_L^3 + 3B_R^3) + (2B_{H_1}^3 + 2B_{H_2}^3 + B_S^3) \\ &= -3B_{H_1}^3 - 3(3B_L + 18B_Q - 7B_S)B_{H_1}^2 - 3(3B_L^2 + (18B_Q - 7B_S)B_S)B_{H_1} + 3B_L^3 + B_S(27B_Q^2 - 27B_S B_Q + 8B_S^2) \\ \mathcal{A}_{BYY} &= 3(6B_Q Y_Q^2 + 3B_{U_R} Y_{U_R}^2 + 3B_{D_R} Y_{D_R}^2) + (6B_L Y_L^2 + 3B_R Y_R^2) + (2B_{H_1} + 2B_{H_2})Y_{H_1}^2 \\ &= \frac{1}{2}(-3B_L - 9B_Q + 7B_S) \\ \mathcal{A}_{BBY} &= 3(6B_Q^2 Y_Q + 3B_{U_R}^2 Y_{U_R} + 3B_{D_R}^2 Y_{D_R}) + (6B_L^2 Y_L + 3B_R^2 Y_R) + (2B_{H_1}^2 - 2B_{H_2}^2)Y_{H_1} \\ &= 2B_{H_1}(3B_L + 9B_Q - 5B_S) + (12B_Q - 5B_S)B_S \\ \mathcal{A}_{BWW} &= \frac{1}{2}(18B_Q + 6B_L + 2B_{H_1} + 2B_{H_2}) = 3B_L + 9B_Q - B_S \\ \mathcal{A}_{BGG} &= \frac{1}{2}(6B_Q + 3B_{U_R} + 3B_{D_R}) = \frac{3}{2}B_S,\end{aligned}\quad (4)$$

where Y_Q, Y_L are the hypercharges of the left-handed doublets of the quarks and leptons, respectively, while Y_{U_R}, Y_{D_R}, Y_R are the hypercharges of the $\hat{U}_R, \hat{D}_R, \hat{R}$ superfields which correspond to the hypercharges of the right-handed quarks and leptons, with the opposite sign.

In the absence of a specific charge assignment coming from a string (or other) construction, these equations can be interpreted as defining conditions of a specific model. The role of string theory or of any other construction is to fix the charges, but for the rest the basic structure remains determined by the approach outlined below, and as such is truly general.

V. THE STÜCKELBERG MULTIPLER

In supersymmetric models the cancellation of the anomaly using the Wess-Zumino (WZ) counterterm can be obtained by the introduction of a Stückelberg supermultiplet, associated with the extra $U(1)$. The multiplet contributes to the supersymmetric version of the Stückelberg mechanism [14] and in the WZ interaction that describes the coupling of the supermultiplet to the gauge supermultiplet. We recall that in anomaly-free theories the Stückelberg mechanism has the feature of contributing to the mass of the anomalous gauge boson, eventually also in combination with the Higgs sector [25,26]. This construction holds both in the nonsupersymmetric and in the supersymmetric case.

Obviously, the presence of a mixing between the Higgs and Stückelberg components in the potential of more ge-

neric models in an anomaly-free theory, produces a new CP -odd component in the scalar sector, but deprived of axionlike couplings. On the contrary, these couplings appear in the case in which the two mechanisms (the Higgs and the Stückelberg) involve an anomalous $U(1)$, due to the presence of Wess-Zumino terms, for specific superpotentials. These interactions are induced in the effective action by the mechanism of anomaly cancellation.

The Lagrangian describing the Stückelberg supermultiplet is given by [14]

$$\mathcal{L}_{\text{st}} = \int d^4\theta [2M_{\text{st}}\hat{B} + \hat{\mathbf{b}} + \hat{\mathbf{b}}^\dagger]^2 \quad (5)$$

where \hat{B} is the Abelian scalar superfield associated to the extra $U(1)_B$, $\hat{\mathbf{b}}$ is a left-chiral superfield and M_{st} is the Stückelberg mass.

The former Lagrangian is invariant under the following gauge transformations

$$\hat{B} \rightarrow \hat{B}' + i(\hat{\Lambda} - \hat{\Lambda}^\dagger) \quad \hat{\mathbf{b}} \rightarrow \hat{\mathbf{b}}' - i2M_{\text{st}}\hat{\Lambda} \quad (6)$$

where $\hat{\Lambda}$ is a generic left-chiral superfield. Introducing the component fields expansion we obtain

$$\hat{B} = -\theta\sigma^\mu\bar{\theta}B_\mu + i\theta\theta\bar{\theta}\bar{\lambda}_B - i\bar{\theta}\bar{\theta}\theta\lambda_B + \frac{1}{2}\theta\theta\bar{\theta}\bar{\theta}D_B \quad (7)$$

$$\begin{aligned} \hat{\mathbf{b}} &= b + i\sqrt{2}\theta\psi_{\mathbf{b}} - i\theta\sigma^\mu\bar{\theta}\partial_\mu b + \frac{\sqrt{2}}{2}\theta\theta\bar{\theta}\bar{\sigma}^\mu\partial_\mu\psi_{\mathbf{b}} \\ &\quad - \frac{1}{4}\theta\theta\bar{\theta}\bar{\theta}\square b + \theta\theta F_{\mathbf{b}}, \end{aligned} \quad (8)$$

where B_μ is the Stückelberg field, $\lambda_B, \bar{\lambda}_B$ are, respectively, the left- and right-handed Stückelberg gauginos, D_B is the corresponding D -term for the gauge supermultiplet of B_μ , b is a complex scalar field, $\psi_{\mathbf{b}}$ is the supersymmetric axion (axino) and $F_{\mathbf{b}}$ is the F -term of $\hat{\mathbf{b}}$.

After the integration over the Grassman variables the Lagrangian density is given by

$$\begin{aligned} \mathcal{L}_{\text{st}} &= 2(\partial_\mu \text{Im}b + M_{\text{st}}B_\mu)^2 + i\psi_{\mathbf{b}}\sigma^\mu\partial_\mu\bar{\psi}_{\mathbf{b}} \\ &\quad + i\bar{\psi}_{\mathbf{b}}\bar{\sigma}^\mu\partial_\mu\psi_{\mathbf{b}} + 2F_{\mathbf{b}}F_{\mathbf{b}}^\dagger + 4M_{\text{st}}\text{Re}bD_B \\ &\quad - 2\sqrt{2}M_{\text{st}}(\psi_{\mathbf{b}}\lambda_B + \text{H.c.}), \end{aligned} \quad (9)$$

where the auxiliary fields $F_{\mathbf{b}}$ and D_B will be defined in the next sections.

A. The axion Lagrangian

The axion Lagrangian contains the Stückelberg gauge-invariant terms introduced above and the Wess-Zumino interactions for the anomaly cancellation and it is given by

$$\begin{aligned} \mathcal{L}_{\text{axion}} &= \frac{1}{2}(\partial_\mu \text{Im}b + M_{\text{st}}B_\mu)^2 + \frac{i}{4}\psi_{\mathbf{b}}\sigma^\mu\partial_\mu\bar{\psi}_{\mathbf{b}} + \frac{i}{4}\bar{\psi}_{\mathbf{b}}\bar{\sigma}^\mu\partial_\mu\psi_{\mathbf{b}} + \frac{1}{2}F_{\mathbf{b}}F_{\mathbf{b}}^\dagger - \frac{M_{\text{st}}}{\sqrt{2}}(\psi_{\mathbf{b}}\lambda_B + \text{H.c.}) \\ &\quad - \frac{1}{16}b_G\epsilon^{\mu\nu\rho\sigma}G_{\mu\nu}^a G_{\rho\sigma}^a \text{Im}b - \frac{1}{16}b_W\epsilon^{\mu\nu\rho\sigma}W_{\mu\nu}^i W_{\rho\sigma}^i \text{Im}b - \frac{1}{4}b_Y\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}^Y F_{\rho\sigma}^Y \text{Im}b - \frac{1}{4}b_B\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}^B F_{\rho\sigma}^B \text{Im}b \\ &\quad - \frac{1}{4}b_{YB}\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}^Y F_{\rho\sigma}^B \text{Im}b + \frac{1}{4}b_W\left[\frac{1}{4}\text{Im}b\lambda_{W^i}\sigma^\mu D_\mu\bar{\lambda}_{W^i} - \frac{i}{4\sqrt{2}}\psi_{\mathbf{b}}\lambda_{W^i}\sigma^\mu\bar{\sigma}^\nu W_{\mu\nu}^i + \frac{1}{4}F_{\mathbf{b}}\lambda_{W^i}\lambda_{W^i}\right. \\ &\quad \left. + \frac{1}{2\sqrt{2}}\psi_{\mathbf{b}}\lambda_{W^i}D^i + \text{H.c.}\right] + \frac{1}{4}b_G\left[\frac{1}{4}\text{Im}b\lambda_{g^a}\sigma^\mu D_\mu\bar{\lambda}_{g^a} - \frac{i}{4\sqrt{2}}\psi_{\mathbf{b}}\lambda_{g^a}\sigma^\mu\bar{\sigma}^\nu G_{\mu\nu}^a + \frac{1}{4}F_{\mathbf{b}}\lambda_{g^a}\lambda_{g^a}\right. \\ &\quad \left. + \frac{1}{2\sqrt{2}}\psi_{\mathbf{b}}\lambda_{g^a}D^a + \text{H.c.}\right] + b_Y\left[\text{Im}b\lambda_Y\sigma^\mu D_\mu\bar{\lambda}_Y - \frac{i}{2\sqrt{2}}\psi_{\mathbf{b}}\lambda_Y\sigma^\mu\bar{\sigma}^\nu F_{\mu\nu}^Y + \frac{1}{2}F_{\mathbf{b}}\lambda_Y\lambda_Y + \frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_Y D_Y + \text{H.c.}\right] \\ &\quad + b_B\left[\text{Im}b\lambda_B\sigma^\mu D_\mu\bar{\lambda}_B - \frac{i}{2\sqrt{2}}\psi_{\mathbf{b}}\lambda_B\sigma^\mu\bar{\sigma}^\nu F_{\mu\nu}^B + \frac{1}{2}F_{\mathbf{b}}\lambda_B\lambda_B + \frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_B D_B + \text{H.c.}\right] + b_{YB}\left[\left(\text{Im}b\lambda_Y\sigma^\mu\partial_\mu\bar{\lambda}_B\right.\right. \\ &\quad \left.\left. + \frac{1}{2}F_{\mathbf{b}}\lambda_Y\lambda_B + \frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_Y D_B - \frac{i}{2\sqrt{2}}\lambda_Y\sigma^\mu\bar{\sigma}^\nu F_{\mu\nu}^B\psi_{\mathbf{b}}\right) + (Y \leftrightarrow B) + \text{H.c.}\right], \end{aligned} \quad (11)$$

with the F and D terms given by

$$\begin{aligned} F_{\mathbf{b}} &= -(b_G\bar{\lambda}_{g^a}\bar{\lambda}_{g^a} + b_W\bar{\lambda}_{W^i}\bar{\lambda}_{W^i} + b_Y\bar{\lambda}_Y\bar{\lambda}_Y + b_B\bar{\lambda}_B\bar{\lambda}_B + b_{YB}\bar{\lambda}_Y\bar{\lambda}_B), \\ D_B &= -\left[\frac{g_B}{2\sqrt{2}}(B_L\tilde{L}^\dagger\tilde{L} + B_R\tilde{R}^\dagger\tilde{R} + B_Q\tilde{Q}^\dagger\tilde{Q} + B_U\tilde{U}_R^\dagger\tilde{U}_R + B_D\tilde{D}_R^\dagger\tilde{D}_R + B_{H_1}H_1^\dagger H_1 + B_{H_2}H_2^\dagger H_2 + B_S S^\dagger S) + \frac{1}{2}\psi_{\mathbf{b}}(b_B\lambda_B + b_{YB}\lambda_Y)\right], \\ D_Y &= -\left[\frac{g_Y}{2\sqrt{2}}(\tilde{L}^\dagger\tilde{L} - 2\tilde{R}^\dagger\tilde{R} - \frac{1}{3}\tilde{Q}^\dagger\tilde{Q} + \frac{4}{3}\tilde{U}_R^\dagger\tilde{U}_R - \frac{2}{3}\tilde{D}_R^\dagger\tilde{D}_R + H_1^\dagger H_1 - H_2^\dagger H_2) + \frac{1}{2}\psi_{\mathbf{b}}(b_Y\lambda_Y + b_{YB}\lambda_B)\right] \\ D^i &= -\frac{1}{2}\left[g_2(\tilde{L}^\dagger\tau^i\tilde{L} + \tilde{Q}^\dagger\tau^i\tilde{Q} + H_1^\dagger\tau^i H_1 + H_2^\dagger\tau^i H_2) + \frac{b_W}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_{W^i}\right] \\ D^a &= -\frac{1}{2}\left[g_s(\tilde{Q}^\dagger T^a\tilde{Q} + \tilde{U}_R^\dagger T^a\tilde{U}_R + \tilde{D}_R^\dagger T^a\tilde{D}_R) + \frac{b_G}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_{g^a}\right], \end{aligned} \quad (12)$$

$$\begin{aligned} \mathcal{L}_{\text{axion}} &= \frac{1}{4}\int d^4\theta(\hat{\mathbf{b}} + \hat{\mathbf{b}}^\dagger + 2M_{\text{st}}\hat{B})^2 - \frac{1}{2}\int d^4\theta\left\{\left[\frac{1}{2}b_G\right.\right. \\ &\quad \times \text{Tr}(\mathcal{G}\mathcal{G})\hat{\mathbf{b}} + \frac{1}{2}b_W\text{Tr}(WW)\hat{\mathbf{b}} + b_Y\hat{\mathbf{b}}W_\alpha^Y W^{Y,\alpha} \\ &\quad \left.\left.+ b_B\hat{\mathbf{b}}W_\alpha^B W^{B,\alpha} + b_{YB}\hat{\mathbf{b}}W_\alpha^Y W^{B,\alpha}\right]\delta(\bar{\theta}^2) + \text{H.c.}\right\}, \end{aligned} \quad (10)$$

where we have denoted with \mathcal{G} the field strength of $SU(3)_c$, with W the supersymmetric field-strength of $SU(2)$, with W^Y and with W^B the supersymmetric field strength of $U(1)_Y$ and $U(1)_B$, respectively. The factors in front of the WZ counterterms (b_X) are determined by the standard conditions of anomaly cancellation. The Lagrangian, in our case, contains extra WZ counterterms with respect to [10], in particular, we need to impose the cancellation of the mixed $B - SU(3) - SU(3)$ anomaly, which is now nonvanishing due to the charges of the two Higgsinos in the model, which are not opposite. In the MSSM this cancellation is identical, due to the specific color charges of the fermions in each generation. This implies that in our case the effective action contains both a bGG interaction of the axion with the gluons and a vertex involving the corresponding gauginos (gluinos).

Expanding the $\mathcal{L}_{\text{axion}}$ in the component fields we obtain

in which we have terms coming both from $\mathcal{L}_{\text{axion}}$ and from the USSM Lagrangian that can be found in the appendix.

B. The kinetic mixing

In these type of supersymmetric models the extra $U(1)_B$ sector can mix with $U(1)_Y$ in different ways. In particular, in the context of USSM-A, the kinetic mixing is treated as in the NMSSM with the inclusion of an anomalous $U(1)_B$ symmetry and the extra singlet \hat{S} is charged under B .

The Lagrangian for the gauge fields is modified by introducing a mixing term $B - Y$ proportional to a small parameter sina

$$\mathcal{L}_{\text{mixing}} = -\frac{1}{4} \int d^4\theta \, 2 \, \text{sina} \, W^{Y\alpha} W_\alpha^B \delta^2(\bar{\theta}) + \text{H.c.} \quad (13)$$

where sina represents the mixing between the two Abelian structures $U(1)_Y$ and $U(1)_B$. In the same way, the gauge mass terms Lagrangian in the presence of kinetic mixing is modified by the inclusion of a term proportional to the mass parameter M_{YB} as follows

$$\mathcal{L}_{\text{GMTmix}} = \frac{1}{2} \int d^4\theta [M_{YB} W^{Y\alpha} W_\alpha^B + \text{H.c.}] \delta^4(\theta, \bar{\theta}). \quad (14)$$

Furthermore, the USSM-A is affected by another source of kinetic mixing coming from the mixed counterterm proportional to b_{YB} in the expression of $\mathcal{L}_{\text{axion}}$. Expanding this expression in component fields we observe that the multiplet $\hat{\mathbf{b}}$ contains the complex scalar field b whose real part can be $\text{Re}b \neq 0$ and it generates a kinetic mixing proportional to $\propto b_{YB} \text{Re}b g_Y g_B$, where the coefficient b_{YB} fixed by the anomaly cancellation procedure, goes like the in-

verse of the Stückelberg mass and can be neglected in this first analysis (see Ref. [10]). In our formulation we assume $\text{sina} = 0$ for simplicity and we will give a more detailed analysis of the kinetic mixing in the context of the USSM-A in a forthcoming paper [27].

C. The Fayet-Iliopoulos terms

To be as more general as possible, in theories with $U(1)$ s gauge superfields we should add to the Lagrangian the following Fayet-Iliopoulos (FI) term

$$\mathcal{L}_{\text{FI}} = \xi_Y D_Y + \xi_B D_B. \quad (15)$$

which is allowed by symmetry reasons. Here ξ_Y , ξ_B are two coefficients, while D_Y and D_B are the D -terms corresponding to the $U(1)_Y$ and $U(1)_B$ symmetry, respectively. In our analysis we omit these contributions even if a quadratically divergent FI always appears in a field theory at one loop [28]. The reason resides in the fact that, in the low-energy Lagrangian there should be a counterterm, which compensates precisely both the divergent and the finite part of the one-loop contributions (see Ref. [10]). We are also omitting the terms responsible for the cancellation of gravitational anomalies. A more comprehensive description will be given in [27].

Some of the notations used in our analysis are recalled in the appendix, here we just mention that the scalars of the model are denoted, as usual, by a tilde ($\tilde{}$). It is convenient to combine the axion sector and the F and D terms extracted from the other sectors of the total Lagrangian of the model. This combination is in general defined to be the auxiliary Lagrangian, or \mathcal{L}_{aux} , which is given by

$$\begin{aligned} \mathcal{L}_{\text{aux}} = & -y_e^2 H_1^\dagger H_1 \tilde{R} \tilde{R}^\dagger - y_u^2 H_2^\dagger H_2 \tilde{U}_R \tilde{U}_R^\dagger - y_d^2 H_1^\dagger H_1 \tilde{D}_R \tilde{D}_R^\dagger - |\lambda H_1 \cdot H_2|^2 - |\lambda S|^2 (H_2^\dagger H_2 + H_1^\dagger H_1) - y_d^2 \tilde{D}_R^\dagger \tilde{D}_R \tilde{Q}^\dagger \tilde{Q} \\ & - y_e^2 \tilde{L}^\dagger \tilde{L} \tilde{R} \tilde{R}^\dagger - y_u^2 \tilde{U}_R \tilde{U}_R^\dagger \tilde{Q}^\dagger \tilde{Q} - \lambda y_u (S \tilde{Q}^\dagger H_1 \tilde{U}_R^\dagger + \text{H.c.}) - \lambda y_d (S \tilde{Q}^\dagger H_2 \tilde{D}_R^\dagger + \text{H.c.}) \lambda y_e (S \tilde{L}^\dagger H_2 \tilde{R}^\dagger + \text{H.c.}) \\ & - y_d y_u (\tilde{U}_R^\dagger H_2^\dagger H_1 \tilde{D}_R + \text{H.c.}) - y_e y_d (\tilde{D}_R^\dagger \tilde{Q}^\dagger \tilde{L} \tilde{R} + \text{H.c.}) + |(b_G \lambda_{g^a} \lambda_{g^a} + b_W \lambda_{W^i} \lambda_{W^i} + b_Y \lambda_Y \lambda_Y + b_B \lambda_B \lambda_B \\ & + b_{YB} \lambda_Y \lambda_B)|^2 - \frac{1}{2} \left[g_s (\tilde{Q}^\dagger T^a \tilde{Q} + \tilde{U}_R^\dagger T^a \tilde{U}_R + \tilde{D}_R^\dagger T^a \tilde{D}_R) + \frac{b_G}{\sqrt{2}} \psi_{\mathbf{b}} \lambda_{g^a} \right]^2 - \frac{1}{2} \left[g_2 (\tilde{L}^\dagger \tau^i \tilde{L} + \tilde{Q}^\dagger \tau^i \tilde{Q} + H_1^\dagger \tau^i H_1 \right. \\ & + H_2^\dagger \tau^i H_2) + \frac{b_W}{\sqrt{2}} \psi_{\mathbf{b}} \lambda_{W^i} \left. \right]^2 - \left[\frac{g_Y}{2\sqrt{2}} (\tilde{L}^\dagger \tilde{L} - 2\tilde{R}^\dagger \tilde{R} - \frac{1}{3} \tilde{Q}^\dagger \tilde{Q} + \frac{4}{3} \tilde{U}_R^\dagger \tilde{U}_R - \frac{2}{3} \tilde{D}_R^\dagger \tilde{D}_R + H_1^\dagger H_1 - H_2^\dagger H_2) \right. \\ & + \frac{1}{2} \psi_{\mathbf{b}} (b_Y \lambda_Y + b_{YB} \lambda_B) \left. \right]^2 - \left[\frac{g_B}{2\sqrt{2}} (B_L \tilde{L}^\dagger \tilde{L} + B_R \tilde{R}^\dagger \tilde{R} + B_Q \tilde{Q}^\dagger \tilde{Q} + B_U \tilde{U}_R^\dagger \tilde{U}_R + B_D \tilde{D}_R^\dagger \tilde{D}_R + B_{H_1} H_1^\dagger H_1 \right. \\ & + B_{H_2} H_2^\dagger H_2 + B_S S^\dagger S) + \frac{1}{2} \psi_{\mathbf{b}} (b_B \lambda_B + b_{YB} \lambda_Y) \left. \right]^2 + \frac{1}{2} [\psi_{\mathbf{b}} \psi_{\mathbf{b}} (b_G^2 \lambda_{g^a} \lambda_{g^a} + b_W^2 \lambda_{W^i} \lambda_{W^i} + (b_Y^2 + b_{YB}^2) \lambda_Y \lambda_Y \\ & + b_B^2 \lambda_B \lambda_B + (b_Y + b_B) b_{YB} \lambda_Y \lambda_B + b_B b_{YB} \lambda_B \lambda_Y) + \text{H.c.}] \end{aligned} \quad (16)$$

where the expressions of the D terms are now determined by Eq. (12).

VI. GOLDSTONES OF THE POTENTIAL AND OF THE MASSIVE GAUGE BOSONS

The identification of the Goldstone modes of the model requires a combined analysis of the potential and of the bilinear mixing terms $Z_i \partial G_{Z_i}$ for all the broken (massive) gauge bosons. Naturally, the expansion near the vacuum is consistent if

the stability conditions of the potential near the expansion point are satisfied. The neutral Goldstone modes corresponding to the physical neutral gauge bosons after the breaking are part of the CP -odd sector together with other physical components, spanning together the entire CP -odd space. In general, in this sector, the potential contains a set of “flat directions,” which appear as Goldstone modes of the matrix of its second derivatives. These Goldstone modes do not necessarily coincide with the Goldstone modes ($G_{Z'}$) identified from the bilinear mixings. This turns out to be the case if the Stückelberg decouples from the scalar potential while it gives mass to one of the anomalous gauge bosons. To clarify this point it is convenient to move back to the nonsupersymmetric case.

The allowed structure of the potential involves b -independent (V) and b -dependent (V') terms, just on the basis of the symmetries of the Lagrangian, given by

$$V = \sum_{a=1,2} (\mu_a^2 H_a^\dagger H_a + \lambda_{aa} (H_a^\dagger H_a)^2) - 2\lambda_{12} (H_1^\dagger H_1)(H_2^\dagger H_2) + 2\lambda'_{12} |H_2^T \tau_2 H_1|^2, \quad (17)$$

and

$$V' = \lambda_0 H_2^\dagger H_1 e^{-i \sum_I (q_2' - q_1')(b_I/M_I)} + \lambda_1 (H_2^\dagger H_1 e^{-i \sum_I (q_2' - q_1')(b_I/M_I)})^2 + \lambda_2 (H_2^\dagger H_2) H_2^\dagger H_1 e^{-i \sum_I (q_2' - q_1')(b_I/M_I)} + \lambda_3 (H_1^\dagger H_1) H_2^\dagger H_1 e^{-i \sum_I (q_2' - q_1')(b_I/M_I)} + c.c. \quad (18)$$

respectively, where the sum over I is a sum over the Stückelberg axions of the (several) anomalous $U(1)$'s. In the supersymmetric case this second contribution is, in general, not allowed, although it might appear after supersymmetry breaking. This second term or “phase-dependent term” is directly responsible for Higgs-axion mixing and for producing a massive axion. The interesting point is that in the supersymmetric case (with b a real field), even if V' is not allowed, we may still, under some particular conditions, end up with a physical axion in the spectrum, as we are now going to elaborate.

As we have mentioned, the identification of the Goldstones of the theory is necessarily done using the kinetic term of the scalars, including the Stückelberg, which in this case takes the form

$$|\mathcal{D}_\mu H_1|^2 + |\mathcal{D}_\mu H_2|^2 + \frac{1}{2}(\partial_\mu b + M_{\text{st}} B_\mu)^2. \quad (19)$$

The expansion of this equation near the stable vacuum gives the usual bilinear mixings characterizing the derivative couplings of the physical massive gauge bosons to the corresponding Goldstones; rather straightforwardly one obtains the combination

$$M_Z Z^\mu \partial_\mu G_Z + M_{Z'} Z'^\mu \partial_\mu G_{Z'} + \dots \quad (20)$$

with G_Z and $G_{Z'}$ being the true Goldstone modes of the theory. Notice, if not obvious, that while G_Z is just expressed as a linear combination of the two CP -odd components of the Higgs, $G_{Z'}$ on the other hand takes a contribution also from b , due to the Stückelberg mass term. Therefore, one of the special features of the combination of the Higgs and Stückelberg mechanisms is that in some cases the potential of the model— V is an example of this situation, since it does not include a b field—is not sufficient to identify all the Goldstone modes. Clearly, if both V and V' are present, then G_Z and $G_{Z'}$ can be extracted from the total potential and coincide with the Goldstone modes extracted from the bilinear mixings of (19) and (20). In this case the physical axion turns out to be massive. We recall that the quadratic part of the CP -odd potential takes the general form

$$V_{CP\text{-odd}} = (\text{Im}H_1^0, \text{Im}H_2^0, b) \mathcal{N} \begin{pmatrix} \text{Im}H_1^0 \\ \text{Im}H_2^0 \\ b \end{pmatrix} \quad (21)$$

for a suitable \mathcal{N} matrix whose explicit expression is important but not necessary for our discussion. In the case of the MSSM the structure of the potential coincides with that of V and one identifies only one physical CP -odd Higgs (called A^0 in the MSSM) which will not have an axionlike coupling, as can be verified by also a simple counting of the degrees of freedom before and after EWSB. In this case the orthogonal transformation that diagonalizes the CP -odd scalar sector takes the form

$$\begin{pmatrix} \text{Im}H_2^0 \\ \text{Im}H_1^0 \end{pmatrix} = O \begin{pmatrix} A^0 \\ G^0 \end{pmatrix} \quad (22)$$

and involves the physical (massive) CP -odd Higgs A^0 and a Goldstone mode G^0 . The above discussion goes through in a similar way also for the anomalous $U(1)$ extension of the MSSM discussed in [10]. For the case of a potential such as $V_{CP\text{-odd}} = V + V'$ instead, there is indeed a mixing between the components of the Higgs and b and the diagonalization of the quadratic part of the potential gives

$$\begin{pmatrix} \text{Im}H_2^0 \\ \text{Im}H_1^0 \\ b \end{pmatrix} = O_3 \begin{pmatrix} \chi \\ G_1^0 \\ G_2^0 \end{pmatrix} \quad (23)$$

with O_3 being an orthogonal matrix. We have denoted the physical field by χ and the NG-bosons by $G_{1,2}^0$. In this case it is rather obvious that χ acquires an axionlike coupling, inherited from b . In other words b has an expansion in terms of χ , G_1^0 and G_2^0 or, equivalently, in terms of χ , G_Z , and $G_{Z'}$, where G_Z and $G_{Z'}$ are identified by Eq. (20). The decomposition is clearly gauge dependent. One important comment concerns the nature of the $bF\tilde{F}$ interactions in this case.

In the unitary gauge the only axionlike couplings left involve the physical component of b , denoted by χ , called “the axi-Higgs,” which gives typical $\chi F\tilde{F}$ interactions. As we have mentioned above, in the absence of V' , b decouples from the rest of the Higgs sector in \mathcal{N} . In this case in the unitary gauge all the anomalous couplings can be removed, and the theory goes back again to its original anomalous form, with the old Lagrangian now replaced by an ordinary *massive* (and possibly anomalous) Yang-Mills theory. It is rather obvious that the truly new element in these types of actions shows up when a physical axionlike particle is induced in the spectrum. In the absence of this, the $bF\tilde{F}$ has dubious meaning, since this term does not cancel the anomaly, as emphasized by Preskill long ago [29]. Rather, it allows a better power-counting of the modified (anomalous) action. A justification of this point of view comes from the fact that an anomalous (and massive) Yang-Mills theory can be given a typical Stückelberg form and a $bF\tilde{F}$ interaction by a field-enlarging transformation [30].

For this reason the only satisfactory potentials are those that either allow b to be part of the scalar sector (such as for $V + V'$) or, alternatively, when they allow, under certain conditions that we are going to discuss next, a mixing between the CP -odd Higgs components and the Stückelberg.

With these motivations in mind, we move to the case of the new superpotential.

VII. SCALAR MASS TERMS, THE SCALAR POTENTIAL AND THE MASS OF THE GAUGE BOSONS

Let us now move to a discussion of the other sectors of the theory, starting from the scalar one. The Lagrangian for

$$\begin{aligned} \mathcal{L}_{\text{aux-F}} = & -y_e^2 H_1^\dagger H_1 \tilde{R} \tilde{R}^\dagger - y_u^2 H_2^\dagger H_2 \tilde{U}_R \tilde{U}_R^\dagger - y_d^2 H_1^\dagger H_1 \tilde{D}_R \tilde{D}_R^\dagger - |\lambda H_1 \cdot H_2|^2 - |\lambda S|^2 (H_2^\dagger H_2 + H_1^\dagger H_1) - y_d^2 \tilde{D}_R^\dagger \tilde{D}_R \tilde{Q}^\dagger \tilde{Q} \\ & - y_e^2 \tilde{L}^\dagger \tilde{L} \tilde{R} \tilde{R}^\dagger - y_u^2 \tilde{U}_R \tilde{U}_R^\dagger \tilde{Q}^\dagger \tilde{Q} - \lambda y_u (S \tilde{Q}^\dagger H_1 \tilde{U}_R^\dagger + \text{H.c.}) - \lambda y_d (S \tilde{Q}^\dagger H_2 \tilde{D}_R^\dagger + \text{H.c.}) \lambda y_e (S \tilde{L}^\dagger H_2 \tilde{R}^\dagger + \text{H.c.}) \\ & - y_d y_u (\tilde{U}_R^\dagger H_2^\dagger H_1 \tilde{D}_R + \text{H.c.}) - y_e y_d (\tilde{D}_R^\dagger \tilde{Q}^\dagger \tilde{L} \tilde{R} + \text{H.c.}), \end{aligned} \quad (27)$$

where the coefficients y_e, y_u, y_d come from the Yukawa couplings of the superpotential, while the D terms are

$$\begin{aligned} \mathcal{L}_{\text{aux-D}} = & -\frac{g_2^2}{2} (\tilde{L}^\dagger \tau^i \tilde{L} + \tilde{Q}^\dagger \tau^i \tilde{Q} + H_1^\dagger \tau^i H_1 + H_2^\dagger \tau^i H_2)^2 - \frac{g_s^2}{2} (\tilde{Q}^\dagger T^a \tilde{Q} + \tilde{U}_R^\dagger T^a \tilde{U}_R + \tilde{D}_R^\dagger T^a \tilde{D}_R)^2 \\ & - \frac{g_Y^2}{8} \left(\tilde{L}^\dagger \tilde{L} - 2\tilde{R}^\dagger \tilde{R} - \frac{1}{3} \tilde{Q}^\dagger \tilde{Q} + \frac{4}{3} \tilde{U}_R^\dagger \tilde{U}_R - \frac{2}{3} \tilde{D}_R^\dagger \tilde{D}_R + H_1^\dagger H_1 - H_2^\dagger H_2 \right)^2 - \frac{g_B^2}{8} (B_L \tilde{L}^\dagger \tilde{L} + B_R \tilde{R}^\dagger \tilde{R} \\ & + B_Q \tilde{Q}^\dagger \tilde{Q} + B_U \tilde{U}_R^\dagger \tilde{U}_R + B_D \tilde{D}_R^\dagger \tilde{D}_R + B_{H_1} H_1^\dagger H_1 + B_{H_2} H_2^\dagger H_2 + B_S S^\dagger S)^2, \end{aligned} \quad (28)$$

where B_L, B_R are the charges of the leptons chiral superfields, B_Q, B_U, B_D are the charges of the left and right chiral superfields of the quark sector and B_{H_1}, B_{H_2}, B_S are the charges of the two Higgs doublet and of the extra singlet, respectively.

the scalar mass terms is given by

$$\begin{aligned} \mathcal{L}_{\text{SMT}} = & -M_L^2 \tilde{L}^\dagger \tilde{L} - m_R^2 \tilde{R}^\dagger \tilde{R} - M_Q^2 \tilde{Q}^\dagger \tilde{Q} - m_{U_R}^2 \tilde{U}_R^\dagger \tilde{U}_R \\ & - m_{D_R}^2 \tilde{D}_R^\dagger \tilde{D}_R - m_1^2 H_1^\dagger H_1 - m_2^2 H_2^\dagger H_2 - m_S^2 S^\dagger S \\ & - (a_\lambda S H_1 \cdot H_2 + \text{H.c.}) - (a_e H_1 \cdot \tilde{L} \tilde{R} + \text{H.c.}) \\ & - (a_d H_1 \cdot \tilde{Q} \tilde{D}_R + \text{H.c.}) - (a_u H_2 \cdot \tilde{Q} \tilde{U}_R + \text{H.c.}), \end{aligned} \quad (24)$$

where $M_L, M_Q, m_R, m_{U_R}, m_{D_R}, m_1, m_2, m_S$ are the mass parameters for the explicit supersymmetry breaking, while a_e, a_λ, a_u, a_d are coefficients with mass dimension one.

The computation of the Lagrangian containing the soft-breaking terms Lagrangian is, as usual, split into the scalar and gaugino mass terms

$$\mathcal{L}_{\text{Soft}} = \mathcal{L}_{\text{SMT}} + \mathcal{L}_{\text{GMT}} + \frac{1}{2} M_{\mathbf{b}} (\psi_{\mathbf{b}} \psi_{\mathbf{b}} + \bar{\psi}_{\mathbf{b}} \bar{\psi}_{\mathbf{b}}), \quad (25)$$

where $M_{\mathbf{b}}$ is a mass parameter for the axino $\psi_{\mathbf{b}}$. The gaugino mass terms given by

$$\begin{aligned} \mathcal{L}_{\text{GMT}} = & -\frac{1}{2} M_G (\lambda_{g^a} \lambda_{g^a} + \bar{\lambda}_{g^a} \bar{\lambda}_{g^a}) - \frac{1}{2} M_w (\lambda_{W^i} \lambda_{W^i} \\ & + \bar{\lambda}_{W^i} \bar{\lambda}_{W^i}) - \frac{1}{2} M_Y (\lambda_Y \lambda_Y + \bar{\lambda}_Y \bar{\lambda}_Y) \\ & - \frac{1}{2} M_B (\lambda_B \lambda_B + \bar{\lambda}_B \bar{\lambda}_B), \end{aligned} \quad (26)$$

where $\lambda_{g^a}, \bar{\lambda}_{g^a}$ are, respectively, the left- and right-handed gauginos of the $SU(3)$ sector, $\lambda_{W^i}, \bar{\lambda}_{W^i}$ are the left- and right-handed gauginos of the $SU(2)$ sector and $\lambda_Y, \bar{\lambda}_Y$ are the chiral gauginos of $U(1)_Y$. The M_G, M_w, M_Y, M_B mass terms are the supersymmetry (SUSY) breaking parameters for $SU(3), SU(2)_w, U(1)_Y$ and $U(1)_B$ respectively. Once we have imposed the equations of motion for the F -terms the on-shell Lagrangian is given by

A. The scalar potential

The study of EWSB in the case of these models proceeds similarly to the USSM [20].

The scalar potential is given by

$$\begin{aligned}
V = & |\lambda H_1 \cdot H_2|^2 + |\lambda S|^2(|H_1|^2 + |H_2|^2) + \frac{1}{8}(g_2^2 + g_Y^2) \\
& \times (H_1^\dagger H_1 - H_2^\dagger H_2)^2 + \frac{g_B^2}{8}(B_{H_1} H_1^\dagger H_1 + B_{H_2} H_2^\dagger H_2 \\
& + B_S S^\dagger S)^2 + \frac{g_2^2}{2}|H_1^\dagger H_2|^2 + m_1^2|H_1|^2 + m_2^2|H_2|^2 \\
& + m_S^2|S|^2 + (a_\lambda S H_1 \cdot H_2 + \text{H.c.}). \quad (29)
\end{aligned}$$

We introduce the following basis

$$\begin{aligned}
H_1 &= \frac{1}{\sqrt{2}} \begin{pmatrix} \text{Re}H_1^0 + i \text{Im}H_1^0 \\ \text{Re}H_1^- + i \text{Im}H_1^- \end{pmatrix}, \\
H_2 &= \frac{1}{\sqrt{2}} \begin{pmatrix} \text{Re}H_2^+ + i \text{Im}H_2^+ \\ \text{Re}H_2^0 + i \text{Im}H_2^0 \end{pmatrix}, \quad (30) \\
S &= \frac{1}{\sqrt{2}} (\text{Re}S + i \text{Im}S),
\end{aligned}$$

where in correspondence of the minimum value of the potential we use the following parametrization for the Higgs fields

$$\langle H_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_1 \\ 0 \end{pmatrix}, \quad \langle H_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 \end{pmatrix}, \quad \langle S \rangle = \frac{v_S}{\sqrt{2}}. \quad (31)$$

As usual, we require the existence of a stable vacuum imposing the conditions

$$\begin{aligned}
m_1^2 v_1 + \frac{1}{2} \lambda^2 v_1 (v_2^2 + v_S^2) + \frac{1}{\sqrt{2}} a_\lambda v_2 v_S - \frac{1}{8} v_1 (v_2^2 - v_1^2) g^2 \\
+ \frac{1}{8} g_B^2 B_{H_1} v_1 (B_{H_2} v_2^2 + B_{H_1} v_1^2 + B_S v_S^2) = 0, \quad (32)
\end{aligned}$$

$$\begin{aligned}
m_2^2 v_2 + \frac{1}{2} \lambda^2 v_2 (v_1^2 + v_S^2) + \frac{1}{\sqrt{2}} a_\lambda v_1 v_S + \frac{1}{8} v_2 (v_2^2 - v_1^2) g^2 \\
+ \frac{1}{8} g_B^2 B_{H_2} v_2 (B_{H_2} v_2^2 + B_{H_1} v_1^2 + B_S v_S^2) = 0, \quad (33)
\end{aligned}$$

$$\begin{aligned}
\frac{1}{\sqrt{2}} a_\lambda v_1 v_2 + m_S^2 v_S + \frac{1}{2} \lambda^2 v_S v^2 + \frac{1}{8} g_B^2 B_S v_S (B_{H_2} v_2^2 \\
+ B_{H_1} v_1^2 + B_S v_S^2) = 0, \quad (34)
\end{aligned}$$

where again a_λ is a mass parameter of the model.

B. Mass of the gauge bosons

The Lagrangian that describes the contributions to the mass of the gauge bosons is given by

$$\begin{aligned}
\mathcal{L}_Q = & |\mathcal{D}_\mu H_1|^2 + |\mathcal{D}_\mu H_2|^2 + |\mathcal{D}_\mu S|^2 \\
& + \frac{1}{2} (\partial_\mu \text{Im}b + M_{\text{st}} B_\mu)^2 \quad (35)
\end{aligned}$$

and involves, beside the two Higgses, the SM bosonic singlet of \hat{S} , the bosonic component of the Stückelberg axion, b , and the Stückelberg mass M_{st} . Collecting the quadratic terms we obtain the contributions to the gauge boson masses which are given by

$$\begin{aligned}
\mathcal{L}_{\text{GM}} = & \frac{g_2^2}{4} (v_1^2 + v_2^2) W^{+\mu} W_\mu^- + \frac{g_2^2}{8} (v_1^2 + v_2^2) W^{3\mu} W_\mu^3 \\
& - \frac{g_2 g_Y}{4} (v_1^2 + v_2^2) W^{3\mu} A_\mu^Y + \frac{g_Y^2}{8} (v_1^2 + v_2^2) A^{Y\mu} A_\mu^Y \\
& + \frac{g_2 g_B}{4} (B_{H_1} v_1^2 - B_{H_2} v_2^2) W_\mu^3 B^\mu - \frac{g_Y g_B}{4} (B_{H_1} v_1^2 \\
& - B_{H_2} v_2^2) A_\mu^Y B^\mu + \frac{g_B^2}{8} (B_{H_1}^2 v_1^2 + B_{H_2}^2 v_2^2 + B_S^2 v_S^2) \\
& \times B_\mu B^\mu + \frac{1}{2} M_{\text{st}}^2 B^\mu B_\mu. \quad (36)
\end{aligned}$$

Using the interaction basis of the gauge field components (W_μ^3, A_μ^Y, B_μ) we obtain the corresponding mass matrix, which is given by

$$\mathcal{M}_{\text{gauge}}^2 = \begin{pmatrix} \frac{g_2^2}{8} v^2 & -\frac{g_2 g_Y}{8} v^2 & \frac{g_2}{8} x_B \\ -\frac{g_2 g_Y}{8} v^2 & \frac{g_Y^2}{8} v^2 & \frac{g_Y}{8} x_B \\ \frac{g_2}{8} x_B & \frac{g_Y}{8} x_B & \frac{N_{BB}}{8} + \frac{M_{\text{st}}^2}{2} \end{pmatrix}, \quad (37)$$

where

$$\begin{aligned}
x_B &= g_B (v_1^2 B_{H_1} - v_2^2 B_{H_2}), \\
N_{BB} &= g_B^2 (B_{H_1}^2 v_1^2 + B_{H_2}^2 v_2^2 + B_S^2 v_S^2), \quad v^2 = v_1^2 + v_2^2. \quad (38)
\end{aligned}$$

Performing the diagonalization we obtain the rotation matrix

$$O_{\text{susy}}^A = \begin{pmatrix} \frac{g_Y}{g} & \frac{g_2}{g} & 0 \\ -\frac{g_2 (f_1 + \sqrt{f_1^2 + 4x_B^2 g^2})}{g \sqrt{2[4g^2 x_B^2 + f_1^2 + f_1 \sqrt{f_1^2 + 4g^2 x_B^2}]}} & \frac{g_Y (f_1 + \sqrt{f_1^2 + 4x_B^2 g^2})}{g \sqrt{2[4g^2 x_B^2 + f_1^2 + f_1 \sqrt{f_1^2 + 4g^2 x_B^2}]}} & \frac{g x_B \sqrt{2}}{\sqrt{[4g^2 x_B^2 + f_1^2 + f_1 \sqrt{f_1^2 + 4g^2 x_B^2}]}} \\ -\frac{g_2 (f_1 - \sqrt{f_1^2 + 4x_B^2 g^2})}{g \sqrt{2[4g^2 x_B^2 + f_1^2 - f_1 \sqrt{f_1^2 + 4g^2 x_B^2}]}} & \frac{g_Y (f_1 - \sqrt{f_1^2 + 4x_B^2 g^2})}{g \sqrt{2[4g^2 x_B^2 + f_1^2 - f_1 \sqrt{f_1^2 + 4g^2 x_B^2}]}} & \frac{g x_B \sqrt{2}}{\sqrt{[4g^2 x_B^2 + f_1^2 - f_1 \sqrt{f_1^2 + 4g^2 x_B^2}]}} \end{pmatrix}, \quad (39)$$

which acts on the interaction basis as

$$\begin{pmatrix} A^\gamma \\ Z \\ Z' \end{pmatrix} = O_{\text{susy}}^A \begin{pmatrix} W^3 \\ A^Y \\ B \end{pmatrix}, \quad (40)$$

and where we have defined $g = \sqrt{g_Y^2 + g_2^2}$ and $f_1 = 4M_{\text{st}}^2 - g^2 v^2 + N_{BB}$.

We obtain one null eigenvalue corresponding to the photon, while the masses of the physical Z and Z' are given by

$$\begin{aligned} M_Z^2 &= \frac{1}{8}(4M_{\text{st}}^2 + g^2 v^2 + N_{BB} - \sqrt{(4M_{\text{st}}^2 - g^2 v^2 + N_{BB})^2 + 4g^2 x_B^2}) \\ M_{Z'}^2 &= \frac{1}{8}(4M_{\text{st}}^2 + g^2 v^2 + N_{BB} + \sqrt{(4M_{\text{st}}^2 - g^2 v^2 + N_{BB})^2 + 4g^2 x_B^2}). \end{aligned} \quad (41)$$

Compared to the nonsupersymmetric case [8], the corrections to the masses of the gauge bosons involve also v_S , which is implicitly contained in N_{BB} .

C. The charged and the CP -even sectors of the scalar potential

The description of the charged sector of the model is performed using the standard basis ($\text{Re}H_2^+$, $\text{Re}H_1^-$). We obtain the following mass matrix

$$\mathcal{M}_c^2 = \begin{pmatrix} \frac{1}{2}(\frac{1}{2}g^2 - \lambda^2)v_1^2 + a_\lambda \frac{v_1 v_S}{\sqrt{2}v_2} & -\frac{1}{2}(\frac{1}{2}g^2 - \lambda^2)v_1 v_2 + a_\lambda \frac{v_S}{\sqrt{2}} \\ -\frac{1}{2}(\frac{1}{2}g^2 - \lambda^2)v_1 v_2 + a_\lambda \frac{v_S}{\sqrt{2}} & \frac{1}{2}(\frac{1}{2}g^2 - \lambda^2)v_2^2 + a_\lambda \frac{v_2 v_S}{\sqrt{2}v_1} \end{pmatrix}. \quad (42)$$

The same mass matrix is obtained in the basis ($-\text{Im}H_2^+$, $\text{Im}H_1^-$). We have one zero eigenvalue corresponding to a charged Goldstone boson and a mass eigenvalue corresponding to the charged Higgs

$$m_{H^\pm}^2 = \left(\frac{v_1}{v_2} + \frac{v_2}{v_1}\right) \left(\frac{1}{4}g^2 v_1 v_2 - \frac{1}{2}\lambda^2 v_1 v_2 + a_\lambda \frac{v_S}{\sqrt{2}}\right). \quad (43)$$

In the analysis of the CP -even sector we use the basis ($\text{Re}H_1^0$, $\text{Re}H_2^0$, $\text{Re}S$). We obtain the matrix elements

$$\begin{aligned} (\mathcal{M}_{\text{ev}}^2)_{11} &= \frac{1}{4}(g_B^2 B_{H_1}^2 + g_Y^2 + g_2^2)v_1^2 - a_\lambda \frac{v_2 v_S}{\sqrt{2}v_1} & (\mathcal{M}_{\text{ev}}^2)_{12} &= \left(\frac{g_B^2}{4} B_{H_1} B_{H_2} + \lambda^2 - \frac{g_2^2 + g_Y^2}{4}\right)v_1 v_2 + a_\lambda \frac{v_S}{\sqrt{2}} \\ (\mathcal{M}_{\text{ev}}^2)_{13} &= a_\lambda \frac{v_2}{\sqrt{2}} + \left(\frac{g_B^2}{4} B_{H_1} B_S + \lambda^2\right)v_1 v_S & (\mathcal{M}_{\text{ev}}^2)_{22} &= \frac{1}{4}(g_B^2 B_{H_1}^2 + g_Y^2 + g_2^2)v_2^2 - a_\lambda \frac{v_2 v_S}{\sqrt{2}v_1} \\ (\mathcal{M}_{\text{ev}}^2)_{23} &= a_\lambda \frac{v_1}{\sqrt{2}} + \left(\frac{g_B^2}{4} B_{H_2} B_S + \lambda^2\right)v_2 v_S & (\mathcal{M}_{\text{ev}}^2)_{33} &= -a_\lambda \frac{v_1 v_2}{\sqrt{2}v_S} + \frac{1}{4}g_B^2 B_S^2 v_S^2 \end{aligned}$$

with the other terms obtained by symmetry ($\mathcal{M}_{12} = \mathcal{M}_{21}$, etc.). The matrix has in general three massive eigenvalues corresponding to the three neutral Higgs particles (H_1^0 , H_2^0 , H_3^0).

D. The neutral CP -odd sector and the axion

The key sector that is responsible for the presence of a physical axion is the CP -odd one. Choosing the basis given by the components ($\text{Im}S$, $\text{Im}H_1^0$, $\text{Im}H_2^0$), our superpotential with an extra singlet gives the mixing matrix

$$\mathcal{M}_{\text{odd}}^2 = \frac{a_\lambda}{\sqrt{2}} \begin{pmatrix} \frac{v_1 v_2}{v_S} & v_2 & v_1 \\ v_2 & \frac{v_2 v_S}{v_1} & v_S \\ v_1 & v_S & \frac{v_1 v_S}{v_2} \end{pmatrix}. \quad (44)$$

Diagonalizing this mass matrix we can identify the orthogonal transformation O^{odd} from the interaction to the mass eigenstates which is given by

$$\begin{pmatrix} \text{Im}S \\ \text{Im}H_1^0 \\ \text{Im}H_2^0 \end{pmatrix} = O^{\text{odd}} \begin{pmatrix} G_1^0 \\ G_2^0 \\ H_4^0 \end{pmatrix}. \quad (45)$$

A simple analysis gives two null eigenvalues, corresponding to two neutral Goldstone bosons, and one physical state, which is identified with a massive neutral Higgs boson

$$m_{H_4^0}^2 = \frac{a_\lambda}{\sqrt{2}} \left(\frac{v_1 v_2}{v_S} + \frac{v_1 v_S}{v_2} + \frac{v_S v_2}{v_1}\right). \quad (46)$$

From the diagonalization procedure we obtain

$$O^{\text{odd}} = \begin{pmatrix} -\frac{v_S}{\sqrt{v_S^2 + v_2^2}} & -\frac{v_S}{\sqrt{v_S^2 + v_1^2}} & \frac{v_1 v_2}{\sqrt{v_1^2 v_2^2 + v^2 v_S^2}} \\ 0 & \frac{v_1}{\sqrt{v_S^2 + v_1^2}} & \frac{v_2 v_S}{\sqrt{v_1^2 v_2^2 + v^2 v_S^2}} \\ \frac{v_2}{\sqrt{v_S^2 + v_2^2}} & 0 & \frac{v_1 v_S}{\sqrt{v_1^2 v_2^2 + v^2 v_S^2}} \end{pmatrix} \quad (47)$$

and the states are given by

$$G_1^0 = \frac{v_2 \text{Im}H_2^0 - v_S \text{Im}S}{\sqrt{v_2^2 + v_S^2}}, \quad G_2^0 = \frac{v_1 \text{Im}H_1^0 - v_S \text{Im}S}{\sqrt{v_1^2 + v_S^2}},$$

$$H_4^0 = \frac{v_1 v_2 \text{Im}S + v_S v_2 \text{Im}H_1^0 + v_1 v_S \text{Im}H_2^0}{\sqrt{v_1^2 v_2^2 + v_S^2 v^2}}, \quad (48)$$

where G_1^0 and G_2^0 are two Goldstone modes, while H_4^0 is the physical Higgs.

Having identified the Goldstones of the potential in the CP -odd sector, the parallel identification of the Goldstones of the massive gauge bosons after EWSB is performed by an analysis of the bilinear mixings. In fact, from the Lagrangian density we can extract the following derivative coupling terms

$$\mathcal{L}_{\text{DC}} = \frac{1}{2}g_2 W_\mu^3 \partial^\mu G_Y - \frac{1}{2}g_Y A_\mu^Y \partial^\mu G_Y + \frac{1}{2}g_B B_\mu \partial^\mu G_B \quad (49)$$

where we have defined

$$G_Y = (v_1 \text{Im}H_1^0 - v_2 \text{Im}H_2^0)$$

$$G_B = (B_{H_1} v_1 \text{Im}H_1^0 + B_{H_2} v_2 \text{Im}H_2^0 + B_S v_S \text{Im}S) + \frac{2M_{\text{st}}}{g_B} \text{Im}b \quad (50)$$

which can be rotated onto the basis (A_μ^Y, Z_μ, Z'_μ) using the O_{susy}^A matrix

$$W_\mu^3 = O_{W\gamma}^A A_\mu^Y + O_{WZ}^A Z_\mu + O_{WZ'}^A Z'_\mu$$

$$A_\mu^Y = O_{Y\gamma}^A A_\mu^Y + O_{YZ}^A Z_\mu + O_{YZ'}^A Z'_\mu \quad (51)$$

$$B_\mu = O_{B\gamma}^A A_\mu^Y + O_{BZ}^A Z_\mu + O_{BZ'}^A Z'_\mu$$

to obtain the expression for \mathcal{L}_{DC} in terms of physical states

$$\mathcal{L}_{\text{DC}} = M_Z Z_\mu \partial^\mu G_Z + M_{Z'} Z'_\mu \partial^\mu G_{Z'}. \quad (52)$$

The two Goldstone modes corresponding to the physical massive gauge bosons are given by

$$M_Z G_Z = -A \left[\frac{v_1}{2x_B} (f_1 + \sqrt{f_1^2 + 4g^2 x_B^2}) - v_1 g_B B_{H_1} \right] \text{Im}H_1^0$$

$$+ A \left[\frac{v_2}{2x_B} (f_1 + \sqrt{f_1^2 + 4g^2 x_B^2}) + v_2 g_B B_{H_2} \right] \text{Im}H_2^0$$

$$+ B_S g_B v_S A \text{Im}S + 2M_{\text{st}} A \text{Im}b \quad (53)$$

$$M_{Z'} G_{Z'} = A' \left[\frac{v_1}{2x_B} (\sqrt{f_1^2 + 4g^2 x_B^2} - f_1) + v_1 g_B B_{H_1} \right] \text{Im}H_1^0$$

$$- A' \left[\frac{v_2}{2x_B} (\sqrt{f_1^2 + 4g^2 x_B^2} - f_1) - v_2 g_B B_{H_2} \right] \text{Im}H_2^0$$

$$+ B_S g_B v_S A' \text{Im}S + 2M_{\text{st}} A' \text{Im}b \quad (54)$$

where we have defined the following coefficients

$$A = \sqrt{\frac{1}{8} - \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, \quad A' = \sqrt{\frac{1}{8} + \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}. \quad (55)$$

It is simple to observe that G_Z and $G_{Z'}$ are orthonormal. At this point, a simple counting of the physical degrees of freedom before and after EWSB can give us a hint on the properties of this model.

Before EWSB we have 10 degrees of freedom: two for A_μ^Y , two for W_μ^3 , three for B_μ , two for the Higgs fields $\text{Im}H_1^0$ and $\text{Im}H_2^0$ and one for the singlet $\text{Im}S$. After the breaking, we are left with two polarization states for the physical photon, 3 degrees of freedom for the Z and the Z' respectively, one neutral Higgs state H_4^0 and one physical state which we are going to identify as the axi-Higgs. Therefore we can build this new physical state requiring its orthogonality with respect to the basis $\{H_4^0, G_Z, G_{Z'}\}$ where H_4^0 , identified as the physical direction of the potential, clearly belongs to the CP -odd sector. We start from the following linear combination

$$\chi = b_1 \text{Im}H_1^0 + b_2 \text{Im}H_2^0 + b_3 \text{Im}S + b_4 \text{Im}b \quad (56)$$

and we determine the coefficients b_1, \dots, b_4 by the following constraints

$$Y_1 = b_3 v_1 v_2 + b_2 v_1 v_S + b_1 v_2 v_S = 0,$$

$$Y_2 = 4b_4 M_{\text{st}} x_B + 2b_3 B_S v_S g_B x_B - b_1 v_1 (f_1 - 2B_{H_1} g_B x_B$$

$$+ \sqrt{f_1^2 + 4g^2 x_B^2}) + b_2 v_2 (f_1 + 2B_{H_2} g_B x_B$$

$$+ \sqrt{f_1^2 + 4g^2 x_B^2}) = 0$$

$$Y_3 = 4b_4 M_{\text{st}} x_B + 2b_3 B_S v_S g_B x_B + b_2 v_2 (f_1 + 2B_{H_2} g_B x_B$$

$$- \sqrt{f_1^2 + 4g^2 x_B^2}) + b_1 v_1 (-f_1 + 2B_{H_1} g_B x_B$$

$$+ \sqrt{f_1^2 + 4g^2 x_B^2}) = 0, \quad (57)$$

which give

$$b_1 = b_4 \frac{2M_{\text{st}}}{g_B B_S} \frac{v_1 v_2^2}{(v_1^2 v_2^2 + v^2 v_S^2)}$$

$$b_2 = b_4 \frac{M_{\text{st}}}{4g_B B_S} \frac{v_1^2 v_2}{(v_1^2 v_2^2 + v^2 v_S^2)}$$

$$b_3 = -b_4 \frac{M_{\text{st}}}{4g_B B_S} \frac{v^2 v_S}{(v_1^2 v_2^2 + v^2 v_S^2)},$$

where the coefficient b_4 is constrained by the normalization of the eigenstates. The physical axion will be given by

$$\begin{aligned} \chi &= \frac{1}{N_\chi} [2M_{\text{st}} v_1 v_2^2 \text{Im}H_1^0 + 2M_{\text{st}} v_1^2 v_2 \text{Im}H_2^0 \\ &\quad - 2M_{\text{st}} v^2 v_S \text{Im}S + B_S g_B (v^2 v_S^2 + v_1^2 v_2^2) \text{Im}b] \\ N_\chi &= \sqrt{4M_{\text{st}}^2 v^2 (v^2 v_S^2 + v_1^2 v_2^2) + B_S^2 g_B^2 (v^2 v_S^2 + v_1^2 v_2^2)^2} \end{aligned} \quad (58)$$

where the new identified state has a nonvanishing projection over the Stückelberg field. Reexpressing $\text{Im}b$ in terms of χ and the Goldstone modes of the massive gauge bosons, we discover that the axionlike interactions (Wess-Zumino terms) mediated by the Stückelberg field can be rotated over χ , giving trilinear vertices of the form $\chi F_I \wedge F_J$, where I and J denote the physical gauge bosons.

The rotation matrix O_{susy}^χ that rotates the physical components and the Goldstones in the CP -odd sector takes the form

$$\begin{pmatrix} H_4^0 \\ G_Z \\ G'_Z \\ \chi \end{pmatrix} = (O_{\text{susy}}^\chi) \begin{pmatrix} \text{Im}H_1^0 \\ \text{Im}H_2^0 \\ \text{Im}S \\ \text{Im}b \end{pmatrix}, \quad (59)$$

where all the entries are defined in Appendix B.

The $B_S = 0$ case: No physical axions

In the case $B_S = 0$, corresponding to a singlet of the entire gauge symmetry, we can proceed in the same way,

$$\mathcal{M}_{\text{odd}}^2 = \begin{pmatrix} -t_S \frac{\sqrt{2}}{v_S} - a_\lambda \frac{v_1 v_2}{\sqrt{2} v_S} & -a_\lambda \frac{v_2}{\sqrt{2}} & -a_\lambda \frac{v_1}{\sqrt{2}} \\ -a_\lambda \frac{v_2}{\sqrt{2}} & -\frac{v_2}{v_1} (m_{12}^2 + a_\lambda \frac{v_S}{\sqrt{2}}) & -m_{12}^2 - a_\lambda \frac{v_S}{\sqrt{2}} \\ -a_\lambda \frac{v_1}{\sqrt{2}} & -m_{12}^2 - a_\lambda \frac{v_S}{\sqrt{2}} & -\frac{v_1}{v_2} (m_{12}^2 + a_\lambda \frac{v_S}{\sqrt{2}}) \end{pmatrix} \quad (61)$$

This sector provides two physical Higgs states and one Goldstone mode of the form²

$$G_{n\text{MSSM}}^0 = \frac{1}{\sqrt{v_1^2 + v_2^2}} (v_1 \text{Im}H_1^0 - v_2 \text{Im}H_2^0). \quad (62)$$

The other Goldstone mode is obtained from the derivative coupling of the Stückelberg term ($B^\mu \partial_\mu \text{Im}b$).

Thus, from the derivative couplings, once we have performed a rotation on the physical basis, we obtain the two orthogonal Goldstone modes $G_Z, G_{Z'}$ corresponding to the Z and the Z' bosons, which are a linear combination of $\text{Im}b$ and of the Goldstone mode obtained from the CP -odd

¹At this stage we do not consider a cubic term in \hat{S} in order to avoid the problem related to the formation of cosmological domain walls (see [22,31,32]), though even in this case one has two Higgs bosons and one Goldstone mode in the CP -odd sector.

²The same Goldstone mode can be obtained from the NMSSM scalar potential [33].

obtaining, however, a different result compared to the previous case. In this case the general structure of the scalar potential can be modified by introducing linear or cubic terms in \hat{S} , corresponding to the same structure of the nMSSM or of the NMSSM, with an additional $U(1)_B$ symmetry. Adding a linear term we obtain¹

$$\begin{aligned} V &= \left| \lambda H_1 \cdot H_2 + \frac{m_{12}^2}{\lambda} \right|^2 + |\lambda S|^2 (|H_1|^2 + |H_2|^2) \\ &\quad + \frac{1}{8} (g_2^2 + g_Y^2) (H_1^\dagger H_1 - H_2^\dagger H_2)^2 \\ &\quad + \frac{g_B^2}{8} B_{H_1}^2 (H_1^\dagger H_1 - H_2^\dagger H_2)^2 + \frac{g_2^2}{2} |H_1^\dagger H_2|^2 + m_1^2 |H_1|^2 \\ &\quad + m_2^2 |H_2|^2 + m_S^2 |S|^2 + (a_\lambda S H_1 \cdot H_2 + t_S S + \text{H.c.}), \end{aligned} \quad (60)$$

where we have introduced the mass parameter m_{12}^2/λ —which is the coefficient of \hat{S} in the nMSSM superpotential—and t_S , which is the coefficient of \hat{S} in the soft-breaking Lagrangian and has mass dimension three. Notice that we have used the condition $B_{H_1} = -B_{H_2}$. In this case, in the basis $\{\text{Im}S, \text{Im}H_1^0, \text{Im}H_2^0\}$, the CP -odd mass matrix is given by

sector,

$$\begin{aligned} G_Z &= \alpha_1 G_{n\text{MSSM}}^0 + \alpha_2 \text{Im}b, \\ G_{Z'} &= \alpha'_1 G_{n\text{MSSM}}^0 + \alpha'_2 \text{Im}b, \end{aligned} \quad (63)$$

where the coefficients $\alpha_1, \dots, \alpha'_2$ are not given in an explicit form for simplicity.

In this case the number of degrees of freedom before the symmetry breaking is again equal to ten. In fact we have two for W_3 , three for B , two for Y , and finally $\text{Im}H_1^0, \text{Im}H_2^0$, and $\text{Im}b$. After EWSB we are left with 3 degrees of freedom for the Z , three for the Z' , two for the photon and two neutral Higgs states, which are physical. Therefore we do not have Higgs-axion mixing.

VIII. THE SFERMION SECTOR

Coming to the scalar fermion sector (sfermions), the Lagrangian in terms of component fields is given by

$$\begin{aligned}
\mathcal{L}_{\text{sfer}}^{\text{MSSM}} = & -\lambda y_e [S^\dagger H_2^\dagger \tilde{L} \tilde{R} + S \tilde{L}^\dagger H_2 \tilde{R}^\dagger] - \lambda y_d [S^\dagger H_2^\dagger \tilde{Q} \tilde{D}_R + S \tilde{Q}^\dagger H_2 \tilde{D}_R^\dagger] - \lambda y_u [S^\dagger H_1^\dagger \tilde{Q} \tilde{U}_R + S \tilde{Q}^\dagger H_1 \tilde{U}_R^\dagger] \\
& - y_e^2 [H_1^\dagger H_1 (\tilde{L}^\dagger \tilde{L} + \tilde{R}^\dagger \tilde{R}) - H_1^\dagger \tilde{L} (H_1^\dagger \tilde{L})^\dagger] - y_d^2 [H_1^\dagger H_1 (\tilde{Q}^\dagger \tilde{Q} + \tilde{D}_R^\dagger \tilde{D}_R) - H_1^\dagger \tilde{Q} (H_1^\dagger \tilde{Q})^\dagger] \\
& - y_u^2 [H_2^\dagger H_2 (\tilde{Q}^\dagger \tilde{Q} + \tilde{U}_R^\dagger \tilde{U}_R) - H_2^\dagger \tilde{Q} (H_2^\dagger \tilde{Q})^\dagger] - M_L^2 \tilde{L}^\dagger \tilde{L} - m_R^2 \tilde{R}^\dagger \tilde{R} - M_Q^2 \tilde{Q}^\dagger \tilde{Q} - m_{U_R}^2 \tilde{U}_R^\dagger \tilde{U}_R - m_{D_R}^2 \tilde{D}_R^\dagger \tilde{D}_R - \\
& \times (a_e H_1 \cdot \tilde{L} \tilde{R} + \text{H.c.}) - (a_d H_1 \cdot \tilde{Q} \tilde{D}_R + \text{H.c.}) - (a_u H_2 \cdot \tilde{Q} \tilde{U}_R + \text{H.c.}) - \frac{g_2^2}{2} (\tilde{L}^\dagger \tau^i \tilde{L} + \tilde{Q}^\dagger \tau^i \tilde{Q} + H_1^\dagger \tau^i H_1 \\
& + H_2^\dagger \tau^i H_2)^2 - \frac{g_3^2}{2} (\tilde{Q}^\dagger T^a \tilde{Q} + \tilde{U}_R^\dagger T^a \tilde{U}_R + \tilde{D}_R^\dagger T^a \tilde{D}_R)^2 - \frac{g_Y^2}{8} \left(\tilde{L}^\dagger \tilde{L} - 2\tilde{R}^\dagger \tilde{R} - \frac{1}{3} \tilde{Q}^\dagger \tilde{Q} + \frac{4}{3} \tilde{U}_R^\dagger \tilde{U}_R - \frac{2}{3} \tilde{D}_R^\dagger \tilde{D}_R \right. \\
& \left. + H_1^\dagger H_1 - H_2^\dagger H_2 \right)^2. \tag{64}
\end{aligned}$$

In the presence of an extra $U(1)_B$ an additional piece coming from the D -terms must be added to the sfermion Lagrangian and it is given by

$$\mathcal{L}_{\text{sfer}}^{U(1)_B} = -\frac{g_B^2}{8} (B_L \tilde{L}^\dagger \tilde{L} + B_R \tilde{R}^\dagger \tilde{R} + B_Q \tilde{Q}^\dagger \tilde{Q} + B_U \tilde{U}_R^\dagger \tilde{U}_R + B_D \tilde{D}_R^\dagger \tilde{D}_R + B_{H_1} H_1^\dagger H_1 + B_{H_2} H_2^\dagger H_2 + B_S S^\dagger S)^2. \tag{65}$$

After spontaneous symmetry breaking we get

$$\begin{aligned}
\mathcal{L}_{\text{sfer}}^{\text{tot}} = & -\frac{1}{2} \lambda v_S y_e v_2 [\tilde{L}^2 \tilde{R} + \tilde{L}^{2\dagger} \tilde{R}^\dagger] - \frac{1}{2} \lambda v_S y_d v_2 [\tilde{Q}^2 \tilde{D}_R + \tilde{Q}^{2\dagger} \tilde{D}_R^\dagger] - \frac{1}{2} \lambda v_S y_u v_1 [\tilde{Q}^1 \tilde{U}_R + \tilde{Q}^{1\dagger} \tilde{U}_R^\dagger] - \frac{1}{2} y_e^2 v_1^2 [\tilde{L}^{2\dagger} \tilde{L}^2 + \tilde{R}^\dagger \tilde{R}] \\
& - \frac{1}{2} y_d^2 v_1^2 [\tilde{Q}^{2\dagger} \tilde{Q}^2 + \tilde{D}_R^\dagger \tilde{D}_R] - \frac{1}{2} y_u^2 v_2^2 [\tilde{Q}^{1\dagger} \tilde{Q}^1 + \tilde{U}_R^\dagger \tilde{U}_R] - M_L^2 \tilde{L}^\dagger \tilde{L} - m_R^2 \tilde{R}^\dagger \tilde{R} - M_Q^2 \tilde{Q}^\dagger \tilde{Q} - m_{U_R}^2 \tilde{U}_R^\dagger \tilde{U}_R - m_{D_R}^2 \tilde{D}_R^\dagger \tilde{D}_R \\
& - \left(a_e \frac{v_1}{\sqrt{2}} \tilde{L}^2 \tilde{R} + \text{H.c.} \right) - \left(a_d \frac{v_1}{\sqrt{2}} \tilde{Q}^2 \tilde{D}_R + \text{H.c.} \right) + \left(a_u \frac{v_2}{\sqrt{2}} \tilde{Q}^1 \tilde{U}_R + \text{H.c.} \right) - \frac{g_2^2}{8} (v_1^2 - v_2^2) (\tilde{L}^{1\dagger} \tilde{L}^1 - \tilde{L}^{2\dagger} \tilde{L}^2 + \tilde{Q}^{1\dagger} \tilde{Q}^1 \\
& - \tilde{Q}^{2\dagger} \tilde{Q}^2) - \frac{g_Y^2}{8} (v_1^2 - v_2^2) \left(\tilde{L}^\dagger \tilde{L} - 2\tilde{R}^\dagger \tilde{R} - \frac{1}{3} \tilde{Q}^\dagger \tilde{Q} + \frac{4}{3} \tilde{U}_R^\dagger \tilde{U}_R - \frac{2}{3} \tilde{D}_R^\dagger \tilde{D}_R \right) - \frac{g_B^2}{8} (B_{H_1} v_1^2 + B_{H_2} v_2^2 + B_S v_3^2) (B_L \tilde{L}^\dagger \tilde{L} \\
& + B_R \tilde{R}^\dagger \tilde{R} + B_Q \tilde{Q}^\dagger \tilde{Q} + B_U \tilde{U}_R^\dagger \tilde{U}_R + B_D \tilde{D}_R^\dagger \tilde{D}_R); \tag{66}
\end{aligned}$$

here and in what follows superscripts on \tilde{L} and \tilde{Q} specify the doublet components. In the basis $(\tilde{L}^2, \tilde{R}^\dagger)$, the entries of the mass matrix are given by

$$\begin{aligned}
(M_{\tilde{L}^2, \tilde{R}^\dagger})_{11} &= y_e^2 \frac{1}{2} v_1^2 + M_L^2 - \frac{1}{8} (g_2^2 - g_Y^2) (v_1^2 - v_2^2) + \frac{g_B^2}{8} B_L (B_{H_1} v_1^2 + B_{H_2} v_2^2 + B_S v_3^2), \\
(M_{\tilde{L}^2, \tilde{R}^\dagger})_{12} &= (M_{\tilde{L}^2, \tilde{R}^\dagger})_{21} = \frac{1}{2} \lambda v_S y_e v_2 + a_e \frac{v_1}{\sqrt{2}}, \\
(M_{\tilde{L}^2, \tilde{R}^\dagger})_{22} &= \frac{1}{2} y_e^2 v_1^2 + m_R^2 - \frac{1}{4} g_Y^2 (v_1^2 - v_2^2) + \frac{g_B^2}{8} B_R (B_{H_1} v_1^2 + B_{H_2} v_2^2 + B_S v_3^2).
\end{aligned} \tag{67}$$

The former matrix can be diagonalized through a rotation defined by

$$\tan 2\theta_{\tilde{L}^2, \tilde{R}^\dagger} = \frac{(\lambda v_S y_e v_2 + a_e \sqrt{2} v_1)}{m_R^2 - M_L^2 + \frac{1}{8} (g_2^2 - 3g_Y^2) (v_1^2 - v_2^2) + \frac{g_B^2}{8} (B_R - B_L) (B_{H_1} v_1^2 + B_{H_2} v_2^2 + B_S v_3^2)}. \tag{68}$$

The eigenvalues have very lengthy expressions and we will omit them for brevity. The three eigenstates are given by

$$\tilde{l}_1 = \cos \theta_{\tilde{L}^2, \tilde{R}^\dagger} \tilde{L}^2 + \sin \theta_{\tilde{L}^2, \tilde{R}^\dagger} \tilde{R}^\dagger \quad \tilde{l}_2 = -\sin \theta_{\tilde{L}^2, \tilde{R}^\dagger} \tilde{L}^2 + \cos \theta_{\tilde{L}^2, \tilde{R}^\dagger} \tilde{R}^\dagger \quad \tilde{l}_3 = \tilde{L}^1. \tag{69}$$

The mass of \tilde{L}^1 is given by

$$M_{\tilde{L}^1}^2 = \frac{1}{8} (g_2^2 + g_Y^2) (v_1^2 - v_2^2) + \frac{g_B^2}{8} B_L (B_{H_1} v_1^2 + B_{H_2} v_2^2 + B_S v_3^2). \tag{70}$$

Using the two basis $(\tilde{Q}^2, \tilde{D}_R^\dagger)$ and $(\tilde{Q}^1, \tilde{U}_R^\dagger)$, the mass sector of the squarks can be written as

$$\mathcal{L}_{\text{squark}} = -(\tilde{Q}^{2\dagger} \quad \tilde{D}_R)M_{\tilde{Q}^2, \tilde{D}_R} \begin{pmatrix} \tilde{Q}^2 \\ \tilde{D}_R^\dagger \end{pmatrix} - (\tilde{Q}^{1\dagger} \quad \tilde{U}_R)M_{\tilde{Q}^1, \tilde{U}_R} \begin{pmatrix} \tilde{Q}^1 \\ \tilde{U}_R^\dagger \end{pmatrix}, \quad (71)$$

where the $M_{\tilde{Q}^2, \tilde{D}_R}$ matrix is defined as

$$\begin{aligned} (M_{\tilde{Q}^2, \tilde{D}_R})_{11} &= \frac{1}{2}y_d^2 v_1^2 + M_Q^2 - \frac{1}{8}\left(g_2^2 + \frac{1}{3}g_Y^2\right)(v_1^2 - v_2^2) + \frac{g_B^2}{8}B_Q(B_{H_1}v_1^2 + B_{H_2}v_2^2 + B_Sv_S^2), \\ (M_{\tilde{Q}^2, \tilde{D}_R})_{12} &= (M_{\tilde{Q}^2, \tilde{D}_R})_{21} = \frac{1}{2}\lambda v_S y_d v_2 + a_d \frac{v_1}{\sqrt{2}}, \\ (M_{\tilde{Q}^2, \tilde{D}_R})_{22} &= \frac{1}{2}y_d^2 v_1^2 + m_{\tilde{D}_R}^2 - \frac{1}{12}g_Y^2(v_1^2 - v_2^2) + \frac{g_B^2}{8}B_{D_R}(B_{H_1}v_1^2 + B_{H_2}v_2^2 + B_Sv_S^2), \end{aligned}$$

while for the $M_{\tilde{Q}^1, \tilde{U}_R}$ matrix we get

$$\begin{aligned} (M_{\tilde{Q}^1, \tilde{U}_R})_{11} &= \frac{1}{2}y_u^2 v_2^2 + M_Q^2 + \frac{1}{8}\left(g_2^2 - \frac{1}{3}g_Y^2\right)(v_1^2 - v_2^2) + \frac{g_B^2}{8}B_Q(B_{H_1}v_1^2 + B_{H_2}v_2^2 + B_Sv_S^2), \\ (M_{\tilde{Q}^1, \tilde{U}_R})_{12} &= (M_{\tilde{Q}^1, \tilde{U}_R})_{21} = \frac{1}{2}\lambda v_S y_u v_1 - a_u \frac{v_2}{\sqrt{2}} \\ (M_{\tilde{Q}^1, \tilde{U}_R})_{22} &= \frac{1}{2}y_u^2 v_2^2 + m_{\tilde{U}_R}^2 + \frac{1}{6}g_Y^2(v_1^2 - v_2^2) + \frac{g_B^2}{8}B_{U_R}(B_{H_1}v_1^2 + B_{H_2}v_2^2 + B_Sv_S^2). \end{aligned} \quad (72)$$

The $M_{\tilde{Q}^2, \tilde{D}_R}$ matrix can be diagonalized using

$$\tilde{q}_1 = \cos\theta_{\tilde{Q}^2, \tilde{D}_R} \tilde{Q}^2 + \sin\theta_{\tilde{Q}^2, \tilde{D}_R} \tilde{D}_R^\dagger \quad \tilde{q}_2 = -\sin\theta_{\tilde{Q}^2, \tilde{D}_R} \tilde{Q}^2 + \cos\theta_{\tilde{Q}^2, \tilde{D}_R} \tilde{D}_R^\dagger,$$

where the $\theta_{\tilde{Q}^2, \tilde{D}_R}$ angle is defined by

$$\tan 2\theta_{\tilde{Q}^2, \tilde{D}_R} = \frac{(\lambda v_S y_d v_2 + a_d \sqrt{2} v_1)}{m_{\tilde{D}_R}^2 - M_Q^2 + \frac{1}{8}(g_2^2 - \frac{1}{3}g_Y^2)(v_1^2 - v_2^2) + \frac{g_B^2}{8}(B_{D_R} - B_Q)(B_{H_1}v_1^2 + B_{H_2}v_2^2 + B_Sv_S^2)}. \quad (73)$$

Again, we omit the explicit expression of the eigenvalues since they are quite lengthy. The $M_{\tilde{Q}^1, \tilde{U}_R}$ matrix can be diagonalized by the following choice

$$\tilde{q}_3 = \cos\theta_{\tilde{Q}^1, \tilde{U}_R} \tilde{Q}^1 + \sin\theta_{\tilde{Q}^1, \tilde{U}_R} \tilde{U}_R^\dagger \quad \tilde{q}_4 = -\sin\theta_{\tilde{Q}^1, \tilde{U}_R} \tilde{Q}^1 + \cos\theta_{\tilde{Q}^1, \tilde{U}_R} \tilde{U}_R^\dagger,$$

where $\theta_{\tilde{Q}^1, \tilde{U}_R}$ is defined by

$$\tan 2\theta_{\tilde{Q}^1, \tilde{U}_R} = \frac{(\lambda v_S y_u \sqrt{2} v_1 - a_u \sqrt{2} v_2)}{m_{\tilde{U}_R}^2 - M_Q^2 - \frac{1}{8}(g_2^2 - \frac{5}{3}g_Y^2)(v_1^2 - v_2^2) + \frac{g_B^2}{8}(B_{U_R} - B_Q)(B_{H_1}v_1^2 + B_{H_2}v_2^2 + B_Sv_S^2)}. \quad (74)$$

Using the parameter values specified in the numerical analysis of the neutralino sector, typical values for sfermion masses are around a few TeV.

IX. WESS-ZUMINO COUNTERTERMS AND CHERN-SIMONS INTERACTIONS

The cancellation of the gauge anomalies in these supersymmetric models are obtained by the introduction of axion counterterms. The supersymmetric form of the corresponding Lagrangian introduces, beside the usual bosonic contributions of the form $bF \wedge F$ additional interactions between the axion and the gauginos and be-

tween the axino, the gauge fields and the corresponding gauginos. It is given by

$$\begin{aligned} \mathcal{L}_C &= - \int d^4\theta \left\{ \left[\frac{1}{2} b_G \text{Tr}(\mathcal{G}\mathcal{G}) \hat{\mathbf{b}} + \frac{1}{2} b_W \text{Tr}(WW) \hat{\mathbf{b}} \right. \right. \\ &\quad + b_Y \hat{\mathbf{b}} W_\alpha^Y W^{Y, \alpha} + b_B \hat{\mathbf{b}} W_\alpha^B W^{B, \alpha} \\ &\quad \left. \left. + b_{YB} \hat{\mathbf{b}} W_\alpha^Y W^{B, \alpha} \right] \delta(\tilde{\theta}^2) + \text{H.c.} \right\}. \end{aligned} \quad (75)$$

whose general e Expanding this expression in component fields using the WZ gauge we obtain

$$\begin{aligned}
\mathcal{L}_C = & -\frac{1}{8}b_G\epsilon^{\mu\nu\rho\sigma}G_{\mu\nu}^aG_{\rho\sigma}^a\text{Im}b - \frac{1}{8}b_W\epsilon^{\mu\nu\rho\sigma}W_{\mu\nu}^iW_{\rho\sigma}^i\text{Im}b - \frac{1}{4}b_Y\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}^YF_{\rho\sigma}^Y\text{Im}b - \frac{1}{4}b_B\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}^BF_{\rho\sigma}^B\text{Im}b \\
& - \frac{1}{4}b_{YB}\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}^YF_{\rho\sigma}^B\text{Im}b + b_G\left[\text{Im}b\frac{1}{2}(\lambda_{g^a}\sigma^\mu D_\mu\bar{\lambda}_{g^a}) - \frac{i}{2\sqrt{2}}\psi_{\mathbf{b}}\frac{1}{2}(\lambda_{g^a}\sigma^\mu\bar{\sigma}^\nu G_{\mu\nu}^a) + \frac{1}{2}F_{\mathbf{b}}\frac{1}{2}(\lambda_{g^a}\lambda_{g^a})\right. \\
& + \left.\frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\frac{1}{2}(\lambda_{g^a}D_G^a) + \text{H.c.}\right] + b_W\left[\text{Im}b\frac{1}{2}(\lambda^a\sigma^\mu D_\mu\bar{\lambda}^a) - \frac{i}{2\sqrt{2}}\psi_{\mathbf{b}}\frac{1}{2}(\lambda_{W^i}\sigma^\mu\bar{\sigma}^\nu W_{\mu\nu}^i) + \frac{1}{2}F_{\mathbf{b}}\frac{1}{2}(\lambda_{W^i}\lambda_{W^i})\right. \\
& + \left.\frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\frac{1}{2}(\lambda_{W^i}D^i) + \text{H.c.}\right] + b_Y\left[\text{Im}b\lambda_Y\sigma^\mu D_\mu\bar{\lambda}_Y - \frac{i}{2\sqrt{2}}\psi_{\mathbf{b}}\lambda_Y\sigma^\mu\bar{\sigma}^\nu F_{\mu\nu}^Y + \frac{1}{2}F_{\mathbf{b}}\lambda_Y\lambda_Y + \frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_Y D_Y + \text{H.c.}\right] \\
& + b_B\left[\text{Im}b\lambda_B\sigma^\mu D_\mu\bar{\lambda}_B - \frac{i}{2\sqrt{2}}\psi_{\mathbf{b}}\lambda_B\sigma^\mu\bar{\sigma}^\nu F_{\mu\nu}^B + \frac{1}{2}F_{\mathbf{b}}\lambda_B\lambda_B + \frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_B D_B + \text{H.c.}\right] \\
& + b_{YB}\left[\left(\text{Im}b\lambda_Y\sigma^\mu\partial_\mu\bar{\lambda}_B + \frac{1}{2}F_{\mathbf{b}}\lambda_Y\lambda_B + \frac{1}{\sqrt{2}}\psi_{\mathbf{b}}\lambda_Y D_B - \frac{i}{2\sqrt{2}}\lambda_Y\sigma^\mu\bar{\sigma}^\nu F_{\mu\nu}^B\psi_{\mathbf{b}}\right) + (Y \leftrightarrow B) + \text{H.c.}\right], \quad (76)
\end{aligned}$$

where we have additional contributions for the cancellation of the $U(1)_B SU(3)SU(3)$ anomaly, which are typical of this model and are not present in previous similar formulations [10].

The Chern-Simons Lagrangian

As we have mentioned above, the Chern-Simons Lagrangian describes the freedom to redistribute the anomaly in the trilinear gauge interactions of AVV and AAA type. In a bottom-up description of these models this freedom is equivalently formulated in terms of external Ward identities on the anomalous vertices. The corresponding Lagrangian is similar to the one given in [10], now with the addition of the gluonic terms. It takes the form

$$\begin{aligned}
\mathcal{L}_{CS} = & -\int d^4\theta\left\{c_1[(\hat{Y}D^\alpha\hat{B} - \hat{B}D^\alpha\hat{Y})W_\alpha^B + \text{H.c.}] - c_2[(\hat{Y}D^\alpha\hat{B} - \hat{B}D^\alpha\hat{Y})W_\alpha^Y + \text{H.c.}] - c_3\text{Tr}[(\hat{W}D^\alpha\hat{B} - \hat{B}D^\alpha\hat{W})W_\alpha\right. \\
& + \left.\frac{1}{6}\hat{W}D^\alpha\hat{B}\hat{D}^2[D_\alpha\hat{W}, \hat{W}] + \text{H.c.}] - c_4\text{Tr}\left[(\hat{G}D^\alpha\hat{B} - \hat{B}D^\alpha\hat{G})\mathcal{G}_\alpha + \frac{1}{6}\hat{G}D^\alpha\hat{B}\hat{D}^2[D_\alpha\hat{G}, \hat{G}] + \text{H.c.}\right]\right\} \quad (77)
\end{aligned}$$

where the coefficients $c_1 \dots c_4$ will be determined by the generalized Ward identities of the model. Expanding this expression in terms of component fields we get

$$\begin{aligned}
\mathcal{L}_{CS} = & -c_1\epsilon^{\mu\nu\rho\sigma}B_\mu Y_\nu F_{\rho\sigma}^B + c_2\epsilon^{\mu\nu\rho\sigma}B_\mu Y_\nu F_{\rho\sigma}^Y + c_3\epsilon^{\mu\nu\rho\sigma}B_\mu\text{Tr}\left(W_\nu F_{\rho\sigma} - \frac{i}{3}W_\nu[W_\rho, W_\sigma]\right) \\
& + c_4\epsilon^{\mu\nu\rho\sigma}B_\mu\text{Tr}\left(G_\nu G_{\rho\sigma} - \frac{i}{3}G_\nu[G_\rho, G_\sigma]\right) - c_1(\lambda_B\sigma^\mu\bar{\lambda}_B A_\mu^Y - \lambda_B\sigma^\mu\bar{\lambda}_Y B_\mu + \text{H.c.}) + c_2(\lambda_Y\sigma^\mu\bar{\lambda}_Y B_\mu \\
& - \lambda_Y\sigma^\mu\bar{\lambda}_B A_\mu^Y + \text{H.c.}) + c_3\text{Tr}(\lambda_W\sigma^\mu\bar{\lambda}_W B_\mu - \lambda_W\sigma^\mu\bar{\lambda}_B W_\mu + \text{H.c.}) + c_4\text{Tr}(\lambda_g\sigma^\mu\bar{\lambda}_g B_\mu - \lambda_g\sigma^\mu\bar{\lambda}_B G_\mu + \text{H.c.}). \quad (78)
\end{aligned}$$

The role of the Lagrangian is to redistribute the anomaly among the three anomalous vertices when the symmetry of the interaction is not enough to fix the partial contributions to the anomaly uniquely.

X. GENERALIZED BROKEN WARD IDENTITIES

The anomaly cancellation mechanism for this supersymmetric model proceeds as in [8,30,34–37], where a detailed description of some physical cases can be found. The resulting anomalies must be cancelled in the Abelian sector BBB , BYY , YBB and in the non-Abelian $SU(2)$ and $SU(3)$ sectors. If we start by using a parametrization of the one-loop trilinear gauge interactions with a symmetric distribution of the AAA anomaly vertex (Δ_{AAA}), in which

we denote with $-k_3 = k_1 + k_2$ the incoming momentum with the λ index and with k_1, k_2 the outgoing momenta, with indices μ and ν respectively, we can introduce generalized Ward identities in the momentum space as defining conditions on the model. We obtain

$$\begin{aligned}
& k_{3,\lambda}\mathcal{A}_{BBB}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) \\
& - \frac{1}{4}b_B\epsilon^{\mu\nu\alpha\beta}k_{1,\alpha}k_{2,\beta} - 2m_f\Delta_{BB}^{\mu\nu} = 0, \quad (79)
\end{aligned}$$

for the BBB case, and analogous conditions in the other sectors. The expressions of Δ_{AAA} , Δ_{BB} and similar are given below; m_f denotes the mass of the fermion in the anomaly loop.

Other two Ward identities are obtained by a cyclic permutation of the momenta. Also, notice that in this

specific case we do not have Chern-Simons interactions in the defining condition. For a BYY triangle we have

$$\begin{aligned} k_{3,\lambda}[\mathcal{A}_{BYY}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) - c_2\varepsilon^{\lambda\mu\nu\alpha}(k_1 - k_2)_\alpha] - \frac{1}{4}b_Y\varepsilon^{\mu\nu\alpha\beta}k_{1,\alpha}k_{2,\beta} - 2m_f\Delta_{YY}^{\mu\nu} &= 0, \\ k_{1,\mu}[\mathcal{A}_{BYY}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) - c_2\varepsilon^{\lambda\mu\nu\alpha}(k_1 - k_2)_\alpha] - 2m_f\Delta_{YY}^{\lambda\nu} &= 0, \\ k_{2,\nu}[\mathcal{A}_{BYY}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) - c_2\varepsilon^{\lambda\mu\nu\alpha}(k_1 - k_2)_\alpha] - 2m_f\Delta_{YY}^{\lambda\mu} &= 0, \end{aligned} \quad (80)$$

where the tensor structure of the triangles is given below. For a YBB triangle we have

$$\begin{aligned} k_{3,\lambda}[\mathcal{A}_{YBB}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) - c_1\varepsilon^{\lambda\mu\nu\alpha}(k_1 - k_2)_\alpha] - 2m_f\Delta_{BB}^{\mu\nu} &= 0, \\ k_{1,\mu}[\mathcal{A}_{YBB}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) - c_1\varepsilon^{\lambda\mu\nu\alpha}(k_1 - k_2)_\alpha] - \frac{1}{4}b_{YB}\varepsilon^{\lambda\nu\alpha\beta}k_{2,\alpha}k_{3,\beta} - 2m_f\Delta_{YB}^{\lambda\nu} &= 0, \\ k_{2,\nu}[\mathcal{A}_{YBB}\Delta_{AAA}^{\lambda\mu\nu}(k_3, k_1, k_2) - c_1\varepsilon^{\lambda\mu\nu\alpha}(k_1 - k_2)_\alpha] - \frac{1}{4}b_{YB}\varepsilon^{\lambda\mu\alpha\beta}k_{3,\alpha}k_{1,\beta} - 2m_f\Delta_{YB}^{\lambda\mu} &= 0, \end{aligned} \quad (81)$$

where the coefficients c_1, c_2 are fixed by the Becchi-Rouet-Stora-Tyutin (BRST) invariance under $U(1)_Y$. The explicit form of the tensors $\Delta_{AAA}^{\lambda\mu\nu}$ and $\Delta_{BB}^{\mu\nu}$, in terms of Feynman integrals, are given by

$$\begin{aligned} \Delta_{AAA}^{\lambda\mu\nu}(m_f \neq 0) &= \frac{1}{\pi^2} \int_0^1 dx \int_0^{1-x} dy \frac{1}{\Delta(m_f)} \left\{ \varepsilon[k_1, \lambda, \mu, \nu] \left[-\frac{\Delta(m_f) - m_f^2}{3} + k_2 \cdot k_2 y(y-1) - xyk_1 \cdot k_2 \right] \right. \\ &+ \varepsilon[k_2, \lambda, \mu, \nu] \left[\frac{\Delta(m_f) - m_f^2}{3} - k_1 \cdot k_1 x(x-1) + xyk_1 \cdot k_2 \right] + \varepsilon[k_1, k_2, \lambda, \nu] (k_1^\mu x(x-1) - xyk_2^\mu) \\ &+ \left. \varepsilon[k_1, k_2, \lambda, \mu] (k_2^\nu y(1-y) + xyk_1^\nu) \right\}, \end{aligned} \quad (82)$$

and

$$\Delta_{BB}^{\mu\nu} = -\frac{m_f}{3\pi^2} \varepsilon^{\mu\nu\alpha\beta} k_{1,\alpha} k_{2,\beta} \int_0^1 \int_0^{1-x} dx dy \frac{1}{\Delta(m_f)}, \quad (83)$$

where $\Delta(m_f) = [m_f^2 + (y-1)yk_2^2 + (x-1)xk_1^2 - 2xyk_1 \cdot k_2]$. For $\Delta_{YY}^{\mu\nu}$ and $\Delta_{YB}^{\mu\nu}$ we obtain similar expressions. The same relations can be reformulated in the mass eigenstate basis in terms of the physical gauge bosons Z and Z' . The structure of the (generalized) Ward identity in this case is shown in Fig. 1, written in configuration space, where the first term corresponds to the anomaly, the second is the axion counterterm projected out on the Goldstone G_Z , and the third diagram describes the mass corrections due to the coupling of the Goldstone to the massive fermion in the loop. In the chiral limit, obviously, the third term is absent.

The generalized Ward identities for the case $U(1)_B SU(2) SU(2)$ have similar expressions, while the case $U(1)_B SU(3) SU(3)$ requires a further comment. As a matter of fact, in this case the Higgsinos do not circulate in the loop, but the BGG triangle exhibits an anomaly when

FIG. 1. The generalized Ward identity for the $Z\gamma\gamma$ vertex in our anomalous model away from the chiral limit. The analogous STI for the SM case consists of only diagrams (a) and (c).

$B_S \neq 0$, [see Eq. (4)]. For the same reason we do not have a BGG anomaly in the MLSOM [8] (minimal low scale orientifold model) case when the Higgs charges under $U(1)_B$ are equal.

XI. Z DECAY INTO FOUR FERMIONS: CHERN-SIMONS INTERACTIONS

One interesting signature of trilinear anomalous vertices involving three anomalous gauge bosons can be investigated in the decay process of the Z/Z' into four fermions by the mediation of two extra anomalous currents. This kind of process is phenomenologically relevant since it is sensitive to the presence of (at least) two or more extra anomalous $U(1)$. As a matter of fact, in the MLSOM (nonsupersymmetric case) in the presence of an Abelian symmetry given by $G_1 = U(1)_Y \times U(1)_B$ where B is anomalous, the off-shell effective vertex does not contain any Chern-Simons interaction by construction. If we take, for instance, the triangle $\langle ZZ'Z' \rangle$, some of the relevant effective vertices coming from the interaction eigenstate basis which have an anomalous component are $\langle BBB \rangle$ and $\langle YBB \rangle$. In the BBB case the Chern-Simons interaction vanishes trivially, while in the YBB case the corresponding Chern-Simons counterterm must be “absorbed” in a redefinition of the triangle in order to ensure the BRST invariance. Equivalently, the YBB vertex does not allow a partial anomaly on the Y leg, since there is no axion for Y . An analysis of the anomalous trilinear interactions in the context of the MLSOM can be found in [35].

In the presence of multiple anomalous $U(1)$'s (such as $U(1)_Y \times U(1)_B \times U(1)_{B'}$) the situation is quite different.

The Z decay into four fermions can be mediated by two different extra neutral currents and the off-shell vertex can be of the type $\langle ZZ'Z'' \rangle$, while from the interaction eigenstate basis a contribution $BB'B'$ appears. A simple inspection of the gauge invariance of this vertex shows that a Chern-Simons interaction can not be absorbed into a redefinition of the $BB'B'$ triangle.

A symmetric distribution of the anomaly on the $BB'B'$ triangle, with outgoing momenta k_1 , k_2 and incoming momentum k , fixes the Rosenberg parametrization as follows³

$$\begin{aligned}
 T_{AAA}^{\lambda\mu\nu} = & \left(-A_5 k_1 \cdot k_2 - A_6 k_2^2 - \frac{a_n}{3} \right) \varepsilon[k_1, \lambda, \mu, \nu] \\
 & + \left(-A_4 k_1 \cdot k_2 - A_3 k_1^2 + \frac{a_n}{3} \right) \varepsilon[k_2, \lambda, \mu, \nu] \\
 & + A_3 k_1^\mu \varepsilon[k_1, k_2, \lambda, \nu] + A_4 k_2^\mu \varepsilon[k_1, k_2, \lambda, \nu] \\
 & + A_5 k_1^\nu \varepsilon[k_1, k_2, \lambda, \mu] + A_6 k_2^\nu \varepsilon[k_1, k_2, \lambda, \mu],
 \end{aligned} \tag{84}$$

thus, we have a partial anomaly equal to $\frac{a_n}{3}$ on each Lorentz index

$$\begin{aligned}
 k^\lambda T_{AAA}^{\lambda\mu\nu} &= \frac{a_n}{3} \varepsilon[k_1, k_2, \mu, \nu] \\
 k_1^\mu T_{AAA}^{\lambda\mu\nu} &= \frac{a_n}{3} \varepsilon[k_1, k_2, \lambda, \nu] \\
 k_2^\nu T_{AAA}^{\lambda\mu\nu} &= -\frac{a_n}{3} \varepsilon[k_1, k_2, \lambda, \mu].
 \end{aligned} \tag{85}$$

The generalized Chern-Simons interaction allowed by the presence of multiple anomalous $U(1)$'s can be formally written as

$$\begin{aligned}
 V_{CS}^{\lambda\mu\nu} = & a_n^{(1)} \varepsilon[\lambda, \mu, \nu, \alpha] (k_1^\alpha - k_2^\alpha) + a_n^{(2)} \varepsilon[\lambda, \mu, \nu, \alpha] \\
 & \times (k_2^\alpha - k_3^\alpha) + a_n^{(3)} \varepsilon[\lambda, \mu, \nu, \alpha] (k_3^\alpha - k_1^\alpha)
 \end{aligned} \tag{86}$$

where $k_3 = -k$ and the coefficients $a_n^{(i)}$ $i = 1, 2, 3$ depend on the model and satisfy the relation $a_n^{(1)} + a_n^{(2)} + a_n^{(3)} = a_n$. Therefore, in the definition of the effective vertex the contributions coming from the Chern-Simons interactions appear explicitly and spoil the symmetric distribution of the anomaly on $BB'B'$. Moreover, the cancellation of the anomaly is ensured by the presence of the WZ interactions, which are constrained by the BRST invariance of the model. For example, the computation of the diagrams described in Figs. 2 and 3 gives

³We have defined $a_n = \frac{i}{2\pi^2}$ and we use the notation $\varepsilon[k_1, k_2, \mu, \nu] = \varepsilon^{\alpha\beta\mu\nu} k_{1,\alpha} k_{2,\beta}$.




FIG. 2. Redefinition of the effective trilinear vertex including the Chern-Simons interactions.




FIG. 3. Decay of the Z boson into 4 fermions plus the Chern-Simons contribution.

$$\begin{aligned}
 \bar{T} = & \varepsilon^\lambda(k) (T_{AAA}^{\lambda\mu\nu} + V_{CS}^{\lambda\mu\nu}) \left[\left(g^{\mu\mu'} - \frac{k_1^\mu k_1^{\mu'}}{M_{Z'}^2} \right) \right. \\
 & \times \frac{-i}{k_1^2 - M_{Z'}^2} \bar{u}(q_1) \Gamma_{\mu'} v(q_2) \left(g^{\nu\nu'} - \frac{k_2^\nu k_2^{\nu'}}{M_{Z''}^2} \right) \\
 & \left. \times \frac{-i}{k_2^2 - M_{Z''}^2} \bar{u}(q_3) \Gamma_{\nu'} v(q_4) \right],
 \end{aligned} \tag{87}$$

where we have indicated with $\Gamma_{\nu'}$ the generic Lorentz structure of the fermion coupling to the extra Z'/Z'' . For instance, the Chern-Simons contribution gives

$$\begin{aligned}
 \bar{T}_{CS} = & \varepsilon^\lambda(k) [a^{(1)} \varepsilon[\lambda, \mu, \nu, k_1 - k_2] + a^{(2)} \varepsilon[\lambda, \mu, \nu, k_2 \\
 & - k_3] + a^{(3)} \varepsilon[\lambda, \mu, \nu, k_3 - k_1]] \\
 & \times \bar{u}(q_1) \Gamma^\mu v(q_2) \bar{u}(q_3) \Gamma^\nu v(q_4) \\
 & \times \frac{-1}{(k_1^2 - M_{Z'}^2)(k_2^2 - M_{Z''}^2)}.
 \end{aligned} \tag{88}$$

The detection of these interactions is rather difficult experimentally, given the low production rates due to the large mass of the extra Z' , currently bound to be larger than 900 GeV.

XII. THE NEUTRALINO SECTOR

Moving to the neutralino sector, here the mass matrix is 7-dimensional because of the presence of the axino, the singlino and the B -ino in the spectrum. In the $B_S \neq 0$ case we obtain

$$\begin{aligned}
\mathcal{L}_{\tilde{\chi}^0} = & -\frac{1}{2}M_w\lambda_{W^3}\lambda_{W^3} - \frac{1}{2}M_Y\lambda_Y\lambda_Y - \frac{1}{2}M_B\lambda_B\lambda_B + \frac{iv_1}{\sqrt{2}}g_2\lambda_{W^3}\tilde{H}_1^1 - \frac{iv_2}{\sqrt{2}}g_2\lambda_{W^3}\tilde{H}_2^2 - \frac{iv_1}{\sqrt{2}}g_Y\lambda_Y\tilde{H}_1^1 + \frac{iv_2}{\sqrt{2}}g_Y\lambda_Y\tilde{H}_2^2 \\
& + \frac{iv_1}{\sqrt{2}}g_B B_{H_1}\lambda_B\tilde{H}_1^1 + \frac{iv_2}{\sqrt{2}}g_B B_{H_2}\lambda_B\tilde{H}_2^2 + \frac{iv_S}{\sqrt{2}}g_B B_S\lambda_B\tilde{S} - \lambda v_S\tilde{H}_1^1\tilde{H}_2^2 - \lambda v_1\tilde{S}\tilde{H}_2^2 - \lambda v_2\tilde{S}\tilde{H}_1^1 + \frac{M_{st}}{2\sqrt{2}}\psi_b\lambda_B \\
& - \frac{1}{2}M_b\psi_b\psi_b + \text{H.c.}, \tag{89}
\end{aligned}$$

where M_w , M_Y , M_B , M_b are mass parameters and the term $\lambda v_S/\sqrt{2}$ plays the role of the μ -term; notice that λ is a dimensionless parameter. We have indicated with λ_{W^3} , λ_Y , λ_B the gauginos of W^3 , A^Y , B respectively and with ψ_b the SUSY particle associated to b . The fields \tilde{H}_1^i and \tilde{H}_2^i ($i = 1, 2$) denote the supersymmetric partners of the two Higgs doublets, while \tilde{S} is the SUSY partner of the extra singlet S .

In the basis $(-i\lambda_{W^3}, -i\lambda_Y, -i\lambda_B, \tilde{H}_1^1, \tilde{H}_2^2, \tilde{S}, -i\psi_b)$ the mass matrix takes the form

$$M_{\tilde{\chi}^0} = \begin{pmatrix} M_w & 0 & 0 & -\frac{v_1}{2}g_2 & \frac{v_2}{2}g_2 & 0 & 0 \\ 0 & M_Y & 0 & \frac{v_1}{2}g_Y & -\frac{v_2}{2}g_Y & 0 & 0 \\ 0 & 0 & M_B & -\frac{v_1}{2}g_B B_{H_1} & -\frac{v_2}{2}g_B B_{H_2} & -\frac{v_S}{2}g_B B_S & -\frac{M_{st}}{\sqrt{2}} \\ -\frac{v_1}{2}g_2 & \frac{v_1}{2}g_Y & -\frac{v_1}{2}g_B B_{H_1} & 0 & -\lambda\frac{v_S}{\sqrt{2}} & -\lambda\frac{v_2}{\sqrt{2}} & 0 \\ \frac{v_2}{2}g_2 & -\frac{v_2}{2}g_Y & -\frac{v_2}{2}g_B B_{H_2} & -\lambda\frac{v_S}{\sqrt{2}} & 0 & -\lambda\frac{v_1}{\sqrt{2}} & 0 \\ 0 & 0 & -\frac{v_S}{2}g_B B_S & -\lambda\frac{v_2}{\sqrt{2}} & -\lambda\frac{v_1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & -\frac{M_{st}}{\sqrt{2}} & 0 & 0 & 0 & M_b \end{pmatrix} \tag{90}$$

that will be analyzed numerically in a section below.

A preliminary choice

A preliminary choice [20] which allows to simplify the structure of the 7×7 neutralino matrix is made by setting $M_w = M_Y = M_B = M_b = \lambda = 0$. In these conditions the diagonalization is rather straightforward and we obtain three null eigenvalues. The first corresponds to a physical pure-photino which is obtained from the rotation

$$\begin{aligned}
\lambda_\gamma &= \sin\theta_W\lambda_{W^3} + \cos\theta_W\lambda_Y, \\
\lambda_{Z_{SM}} &= \cos\theta_W\lambda_{W^3} - \sin\theta_W\lambda_Y, \tag{91}
\end{aligned}$$

where $\lambda_{Z_{SM}}$ is an intermediate unphysical state. The second state, corresponding to a null eigenvalue, is given by a mixture of Higgsino and axino states

$$\tilde{\chi}_2^0 = \frac{M_{st}}{2g_B v_1 B_S}\tilde{H}_1^1 + \frac{M_{st}}{2g_B v_2 B_S}\tilde{H}_2^2 + \psi_b, \tag{92}$$

while the third is a pure Higgsino state which corresponds to the SUSY partner of H_4^0 and it is given by the expression

$$\tilde{\chi}_3^0 = \frac{v_S}{v_1}\tilde{H}_1^1 + \frac{v_S}{v_2}\tilde{H}_2^2 + \tilde{S}. \tag{93}$$

The other states corresponding to the nonzero eigenvalues are complicated combinations of Higgsinos, gauginos ($\lambda_{Z_{SM}}$, λ_B) and the axino.

Notice that in our treatment we are considering for simplicity a real-valued neutralino matrix. In the most general cases—for example in some CP-noninvariant theories—these matrix elements are complex and they may contain phase factors which are physical and can not be eliminated by a redefinition of the fields.

XIII. SUPERSYMMETRIC INTERACTIONS OF THE AXION WITH THE NEUTRALINOS

In this section we proceed with a study of the basic tree-level interaction vertices involving the physical axion (axi-Higgs). Analyzing each sector of the whole Lagrangian we have different types of interactions involving the axi-Higgs.

First of all, from the counterterm Lagrangian we have trilinear interactions obtained by rotating the WZ counterterms on the physical basis, which formally give terms of the type

$$\begin{aligned}
\mathcal{L}_{\chi ZZ} = & R_1\epsilon^{\mu\nu\rho\sigma}Z_{\mu\nu}^{\text{abel}}Z_{\rho\sigma}^{\text{abel}}\chi + R_2\epsilon^{\mu\nu\rho\sigma}Z_{\mu\nu}^{\text{abel}}Z_{\rho\sigma}^{\text{abel}}\chi \\
& + R_3\epsilon^{\mu\nu\rho\sigma}Z_{\mu\nu}^{\text{abel}}Z_{\rho\sigma}^{\text{abel}}\chi, \tag{94}
\end{aligned}$$

where for simplicity we have indicated with R_1 , R_2 , R_3 the coefficients which appear in front of each vertex. These include the rotation matrices, the coupling constants of the gauge groups and the coefficients coming from the anomaly cancellation procedure. We omit their explicit expressions since they are not relevant for this discussion. Notice that in this case only the Abelian part of field strengths contribute to the counterterms for the neutral currents and that $Z_{\mu\nu}^{\text{abel}} = \partial_\mu Z_\nu - \partial_\nu Z_\mu$. The interactions coming from these terms are shown in Fig. 4.

From the axion Lagrangian $\mathcal{L}_{\text{axion}}$ we obtain quadri-linear interactions between χ , the neutralinos/gluinos/charginos, the neutral/charged gauge bosons and trilinear derivative interactions, illustrated in Fig. 5 and 6. In fact, by a careful inspection of $\mathcal{L}_{\text{axion}}$ we find




FIG. 4. Trilinear interactions between χ and the neutral currents.




FIG. 5. Quadrilinear interactions involving χ , charginos/gluinos/neutralinos, and a gauge boson.

$$\begin{aligned} \mathcal{L}_{\text{axion}}^{\chi\tilde{\chi}\tilde{\chi}\text{Gauge}} &= R^Z \chi \tilde{\chi}^\pm \gamma^\mu \tilde{\chi}^\mp Z_\mu + R^G \chi \tilde{G} \gamma^\mu \tilde{G} G_\mu \\ &+ R^W \chi \tilde{\chi}^\pm \Gamma^\mu \tilde{\chi}_i^0 W_\mu^\mp + \{Z \rightarrow Z'\}, \end{aligned} \quad (95)$$

while the derivative trilinear interactions are given by

$$\begin{aligned} \mathcal{L}_{\text{axion}}^{\chi\tilde{\chi}\tilde{G}} &= R^{\chi ij} \chi \tilde{\chi}_i^0 \Gamma^\mu \partial_\mu \tilde{\chi}_j^0 + R^{\chi\tilde{G}} \chi \tilde{G} \gamma^\mu \partial_\mu \tilde{G} \\ &+ R^{\chi\pm} \chi \tilde{\chi}^\pm \Gamma^\mu \partial_\mu \tilde{\chi}^\mp, \end{aligned} \quad (96)$$

where Γ^μ indicates that we can have vector or axial-vector interactions. Trilinear interactions between one neutral current and two axionlike particles can be obtained from $\mathcal{L}_{\text{QUAD}}$ and have the form

$$\begin{aligned} \mathcal{L}_{\text{QUAD}}^{\chi H Z} &= R^{\chi H_i^0 Z} \chi \overleftrightarrow{\partial}^\mu H_i^0 Z_\mu + R^{\chi H^\pm W^\mp} \chi \overleftrightarrow{\partial}^\mu H^\pm W_\mu^\mp \\ &+ \{Z \rightarrow Z'\}; \end{aligned} \quad (97)$$

to these terms correspond the interactions shown in Fig. 7; Analogously, the quadrilinear interactions between two axionlike particles and two neutral gauge bosons are given by (see Fig. 8)




FIG. 6. Derivative trilinear interactions between χ and charginos/gluinos/neutralinos.




FIG. 7. Trilinear interactions between χ , a Higgs boson, and an electroweak gauge boson.

$$\begin{aligned} \mathcal{L}_{\text{QUAD}}^{\chi\chi ZZ} &= R_1^{ZZ} \chi \chi Z_\mu Z^\mu + R_2^{ZZ} \chi H_4^0 Z_\mu Z^\mu \\ &+ R_1^{ZZ'} \chi \chi Z_\mu Z'^\mu + R_2^{ZZ'} \chi H_4^0 Z_\mu Z'^\mu \\ &+ \{Z \rightarrow Z'\} \end{aligned} \quad (98)$$

where, again, we have introduced the coefficients R_i^{ZZ} , R_j^{ZZ} containing the rotation matrices and the couplings, for simplicity.

From the Lagrangian of the scalar mass terms \mathcal{L}_{SMT} we obtain the following trilinear interactions involving the axi-Higgs, the Higgs bosons coming from the scalar sector (CP -even, CP -odd, charged) and the sfermions

$$\begin{aligned} \mathcal{L}_{\text{SMT}}^{\chi\chi\text{even-odd}} &= R^{\chi^2 i} \chi^2 H_i^0 + R^{\chi i} \chi H_4^0 H_i^0 + R^{\chi\pm} \chi H^\mp H^\pm \\ &+ R^{\chi\tilde{f}\tilde{f}} \chi \tilde{f} \tilde{f}, \end{aligned} \quad (99)$$

where H_i^0 with $i = 1, \dots, 3$ indicates the physical Higgs states coming from the CP -even sector (see Fig. 9). We denote with $\mathcal{L}_{\mathcal{W}}$ the on-shell Lagrangian coming from the superpotential, once that the F -terms have been removed, and containing all the Yukawa-type interactions

$$\mathcal{L}_{\mathcal{W}} = \mathcal{L}_{\text{Yuk}} + \mathcal{L}_S + \mathcal{L}_{\text{Yuk-F}} \quad (100)$$

where \mathcal{L}_{Yuk} represents the Yukawa interactions that do not contain the extra singlet S and are linear in y_e, y_u, y_d , while \mathcal{L}_S indicates all the Yukawa interactions containing S . Finally, with $\mathcal{L}_{\text{Yuk-F}}$ we indicate those interactions that are quadratic in y_e, y_u, y_d and in λ . Then we have



FIG. 8. Quadrilinear interaction involving χ , two electroweak neutral gauge bosons, and the CP -odd Higgs.




FIG. 9. Trilinear interaction involving χ and Higgs bosons/sfermions.

$$\begin{aligned}
\mathcal{L}_{\text{Yuk}} = & y_e \epsilon^{ij} [-\tilde{H}_1^i L^j \tilde{R} - \tilde{H}_1^i \tilde{L}^j \tilde{R}^\dagger - H_1^i L^j \tilde{R} - H_1^{i\dagger} \tilde{L}^j R \\
& - \tilde{R} \tilde{H}_1^i \tilde{L}^j - R \tilde{H}_1^i \tilde{L}^j] + y_d \epsilon^{ij} [-\tilde{H}_1^i Q^j \tilde{D}_R \\
& - \tilde{H}_1^i \tilde{Q}^j \tilde{D}_R^\dagger - H_1^i Q^j \tilde{D}_R - H_1^{i\dagger} \tilde{Q}^j D_R - \tilde{D}_R \tilde{H}_1^i \tilde{Q}^j \\
& - D_R \tilde{H}_1^i \tilde{Q}^j] + y_u \epsilon^{ij} [-\tilde{H}_2^i Q^j \tilde{U}_R - \tilde{H}_2^i \tilde{Q}^j \tilde{U}_R^\dagger \\
& - H_2^i Q^j \tilde{U}_R - H_2^{i\dagger} \tilde{Q}^j U_R - \tilde{U}_R \tilde{H}_2^i \tilde{Q}^j - U_R \tilde{H}_2^i \tilde{Q}^j],
\end{aligned} \tag{101}$$

$$\begin{aligned}
\mathcal{L}_S = & -\lambda y_e [S^\dagger H_2^\dagger \tilde{L} \tilde{R} + S \tilde{L}^\dagger H_2 \tilde{R}^\dagger] - \lambda y_d [S^\dagger H_2^\dagger \tilde{Q} \tilde{D}_R \\
& + S \tilde{Q}^\dagger H_2 \tilde{D}_R^\dagger] - \lambda y_u [S^\dagger H_1^\dagger \tilde{Q} \tilde{U}_R + S \tilde{Q}^\dagger H_1 \tilde{U}_R^\dagger] \\
& + \lambda \epsilon^{ij} [-S \tilde{H}_1^i \tilde{H}_2^j - S^\dagger \tilde{H}_1^i \tilde{H}_2^j] \\
& - |\lambda S|^2 (H_2^\dagger H_2 + H_1^\dagger H_1)
\end{aligned} \tag{102}$$

and finally

$$\begin{aligned}
\mathcal{L}_{\text{Yuk-F}} = & -|\lambda H_1 \cdot H_2|^2 - y_e^2 [\tilde{L}^\dagger \tilde{L} \tilde{R}^\dagger \tilde{R} + H_1^\dagger H_1 \\
& \times (\tilde{L}^\dagger \tilde{L} + \tilde{R}^\dagger \tilde{R}) - H_1^\dagger \tilde{L} (H_1^\dagger \tilde{L})^\dagger] \\
& - y_d^2 [\tilde{Q}^\dagger \tilde{Q} \tilde{D}_R^\dagger \tilde{D}_R + H_1^\dagger H_1 (\tilde{Q}^\dagger \tilde{Q} + \tilde{D}_R^\dagger \tilde{D}_R) \\
& - H_1^\dagger \tilde{Q} (H_1^\dagger \tilde{Q})^\dagger] - y_u^2 [\tilde{Q}^\dagger \tilde{Q} \tilde{U}_R^\dagger \tilde{U}_R \\
& + H_2^\dagger H_2 (\tilde{Q}^\dagger \tilde{Q} + \tilde{U}_R^\dagger \tilde{U}_R) - H_2^\dagger \tilde{Q} (H_2^\dagger \tilde{Q})^\dagger].
\end{aligned} \tag{103}$$

From the Yukawa mass terms contained in \mathcal{L}_{Yuk} and in \mathcal{L}_S we can isolate the pseudoscalar coupling of the axi-Higgs to the fermions and a quadrilinear scalar interaction with the sfermions

$$\begin{aligned}
\mathcal{L}_{\text{Yuk-S}}^\chi = & R_{\text{Yuk}}^{\chi \tilde{f} f} \tilde{\psi}_f \gamma^5 \psi_f \chi \\
& + R_S^{\chi^2 \tilde{f} \tilde{f}} \chi \chi \tilde{f} \tilde{f} + R_S^{\chi H_4^0 \tilde{f} \tilde{f}} \chi H_4^0 \tilde{f} \tilde{f}
\end{aligned} \tag{104}$$

where we have indicated with ψ_f the generic fermion and with \tilde{f} the generic sfermion (see Fig. 10).




FIG. 10. Interactions obtained from $\mathcal{L}_{\text{Yuk}}^\chi$.




FIG. 11. Quadrilinear interactions involving χ and CP-odd/CP-even/charged Higgs.

Quadrilinear axionic self interactions can be obtained from \mathcal{L}_S and from $\mathcal{L}_{\text{Yuk-F}}$

$$\begin{aligned}
\mathcal{L}_W^{\chi H_4^0} = & R^{\chi^4} \chi^4 + R^{\chi^3} \chi^3 H_4^0 + R^{\chi^2 \pm} \chi^2 H^\pm H^\mp \\
& + R^{\chi^2} \chi^2 (H_4^0)^2 + R^\chi \chi (H_4^0)^3 + R^{\chi^2 ij} \chi^2 H_i^0 H_j^0 \\
& + R^{\chi H_4^0 ij} \chi H_4^0 H_i^0 H_j^0 + R^{\chi H_4^0 \pm} \chi H_4^0 H^\pm H^\mp
\end{aligned} \tag{105}$$

and are listed in Fig. 11.

XIV. NUMERICAL ANALYSIS

In this section we present a numerical analysis of the neutralino sector. We have performed the numerical diagonalization of the 7×7 neutralino matrix and we have studied the eigenvalues dependence with respect to the free parameters of the model. Furthermore, since in this model the neutralino sector exhibits an axino component due to the presence of Stückelberg interactions, we have investigated, in the case of the lightest neutralino state, its mixing with the other states. In Table II we have listed all the values of the parameters that we have used in our analysis. In our analysis we have followed, in spirit, the approach of Kalinowski and collaborators in [38]. In their paper the authors, who deal with the USSM, present two scenarios: in the first one they assume unified values for the gaugino mass terms and in a second scenario they consider with different values (arbitrary values). We refer to their analysis for further justifications and motivations of this choice. We have chosen $\tan\beta \approx 40$ and we have constrained the value of v_1 in order to be consistent with the value of the mass of the Z_0 boson, while the value of the coupling constant g_B is 0.65.

The values $\lambda < 0.7$ and v_S around 1 TeV are consistent with the MSSM value of the Higgs masses.

The charges B_{H_1} and B_{H_2} are free parameters because we have only four equations coming from the gauge invariance of the superpotential and eight charges to be constrained. One possible choice is $B_{H_1} = -3/(2\sqrt{10})$ and $B_{H_2} = -1/(\sqrt{10})$, which is obtained from the E_6 SSM model [38].

In Figs. 12–15, we plot on the left-hand side the numerical value of the neutralino masses obtained from the diagonalization procedure as a function of the mass parameters $M_{\text{st}}, M_B, M_{\mathbf{b}}, M_Y, M_W$ and of g_B and $\tan\beta$. On the right-hand side we plot the squared value of each component of

TABLE II. Parameters for the neutralino eigenvalues analysis for the charge assignment $B_{H_1} = -3/(2\sqrt{10})$ and $B_{H_2} = -1/(\sqrt{10})$.

	M_Y [TeV]	M_w [TeV]	M_B [TeV]	M_{st} [TeV]	M_b [TeV]	λ	ν_S [TeV]	$\tan\beta$	g_B
Figure 12(a) and 12(b)	1.5	2.5	1.6	0–5	1.5	0.1	0.9	40	0.65
Figure 13(a) and 13(b)	1.5	2.5	0–5	2	1.5	0.1	0.9	40	0.65
Figure 13(c) and 13(d)	1.5	2.5	1.6	2	0–5	0.1	0.9	40	0.65
Figure 14(a) and 14(b)	0–5	2.5	2.1	2	1.5	0.1	0.9	40	0.65
Figure 14(c) and 14(d)	1.5	5–9	2.1	2	1.5	0.1	0.9	40	0.65
Figure 15(a) and 15(b)	1.5	2.5	1.6	2	1.5	0.1	0.9	40	0.1–1
Figure 15(c) and 15(d)	1.5	2.5	1.6	2	1.5	0.1	0.9	1–40	0.65
Figure 15(e) and 15(f)	1.5	2.5	1.6	2.1	1	0.7	0.1–3	40	0.65

the lightest neutralino state in order to establish which component is dominant, since every neutralino state appears as a mixture of the axino, the singlino, etc. We can formally decompose the generic i -th neutralino state ($i = 1, \dots, 7$) in the basis $\{-i\lambda_{W_3}, -i\lambda_Y, -i\lambda_B, \tilde{H}_1^1, \tilde{H}_2^2, \tilde{S}, -i\psi_b\}$


$$\tilde{\chi}_i^0 = a_{i1}\lambda_{W_3} + a_{i2}\lambda_Y + a_{i3}\lambda_B + a_{i4}\tilde{H}_1^1 + a_{i5}\tilde{H}_2^2 + a_{i6}\tilde{S} + a_{i7}\psi_b \quad (106)$$


and in the figures we indicate the square of each component as $c_{ij} = |a_{ij}|^2$, where the lightest state corresponds to the $i = 1$ choice. From the left panel of Figs. 12 and 13 we observe that the value of the mass of the lightest neutralino state that is consistent with the current experimental bounds [39] is obtained approximately by varying the values of M_{st} in the interval 1.7–2.5 TeV, while M_B and M_b in the interval 1–2 TeV. In the right panel of Figs. 12 and 13 it is interesting to observe that for these values of the soft-breaking parameters we have a tiny region beyond 1 TeV in which the axino and the B -ino components are almost coincident, the two Higgsinos are dominant, while the singlino is the most suppressed component. For values of M_{st}, M_B, M_b below 1 TeV and beyond 2.5 TeV, the

lightest neutralino is “mostly” singlino, while the W -ino and the Y -ino components are suppressed and the eigenvalues appear to be nondegenerate apart from the states $\tilde{\chi}_2^0 - \tilde{\chi}_3^0$. From the left-hand side of Fig. 14 it is evident that all the eigenvalues do not exhibit substantial variations with respect to M_Y, M_w and the heaviest states are nondegenerate. In both cases [see Fig. 14(b) and 14(d)], the singlino component is the leading one. A similar feature can be found in the USSM case [38], where the singlino is always dominant with respect to the other components.

Finally, in Fig. 15 we have analyzed the dependence upon the coupling constant $g_B, \tan\beta$ and ν_S . In the left-hand side (a) the mass value of the lightest state starts to be greater than 50 GeV once $g_B > 0.4$ and it is almost degenerate with $\tilde{\chi}_2^0$.

From the analysis of each component in the right panel (b), for g_B less than 0.5 the main contribution comes from the singlino, while the axino and the B -ino are almost degenerate and subdominant with respect to the \tilde{H}_2^2 contribution. When g_B becomes greater than 0.5 we have an inversion: the two Higgsinos are dominant and almost equal, while the singlino is subleading and the combination axino- B -ino is more suppressed.


 FIG. 12 (color online). Study of the neutralino eigenvalues as a function of Stüeckelberg mass M_{st} .


FIG. 13 (color online). The same as Fig. 12 but as a function of M_B and M_b .

As a consequence of our constraint on the vev v_1 , the eigenvalues dependence on $\tan\beta$ is weak [see Fig. 15(c)], while we have a strong impact of low values of $\tan\beta$ on the axino, B -ino and on the singlino components. Even in this case, with the choice of the parameters that we have made in Table II, we can identify a small region in which the contribution of the singlino is highly suppressed.

In the last scenario, represented in Fig. 15(c) and 15(d), it seems possible to have an axino-dominated lightest neutralino. This is achieved with a larger value of the effective μ -term (given by λv_S) and a slightly lower one for the axino SUSY breaking parameter M_b .

Given these results, one important issue that one would like to address concerns the modifications implied by our model respect to standard scenarios of neutralino densities—for instance in the MSSM or in the nMSSM—which require a separate investigation of the (rather large) parameter space. We just remark that a related analysis [40], based on an anomalous version of the MSSM which shares various similarities with our model, shows that for an axino-dominated LSP (light supersymmetric particle)—

in the range between 50 GeV–2 TeV— with a mass gap around 1%–5% between the LSP and the NLSP (next to lightest supersymmetric particle), the constraints from WMAP can be satisfied. The NLSP, in that model, has components which are typical of the (nonanomalous) MSSM, with a dominant gaugino and/or a gaugino-Higgsino projection. In the presence of extra singlets and with a physical axion, which is our case, this scenario should be modified even further, but we expect some similarities with these previous studies, especially in the neutralino sector, to hold. In a recent study of the axion in the MLSOM, for instance, the possibility of having the axion as a long lived particle require a very small mass for this particle ($\sim 10^{-4}$ eV) [41]. In the USSM-A the presence of an axion in the bosonic sector and of a neutralino in the fermionic sector as possible dark-matter components raises the issue of the interplay between the two sectors. At the same time, in the fermionic neutral sector, the role of the coannihilation becomes crucial, especially in the presence of mass degeneracy, which modifies substantially the neutralino relic densities already in this sector. We hope to


 FIG. 14 (color online). The same as Fig. 12 but as a function of M_Y and M_w .

return with a complete analysis of these points in the near future [27]

XV. UNITARITY BOUND OF THE MODEL

Being the theory an effective description of an anomalous Lagrangian in which the presence of the axion is the low-energy signature of a more complicated mechanism of cancellation which would eventually induce higher derivative terms in the effective action, it is necessary at this stage to comment about the unitarity of this class of models. This point has been raised in [36] and further developed in [37]. One of the most natural contexts for discussing unitarity is related to $2 \rightarrow 2$ processes mediated by BIM (Bouchiat—Iliopoulos—Meyer) amplitudes, in particular, those involving gluons and photons. These processes exhibit an anomalous behavior when the $gg \rightarrow \gamma\gamma$ amplitude is mediated by the exchange in the s -channel of neutral gauge bosons that couple to the fermion loops via axial-vector

interactions. As shown in these previous analysis, this class of amplitudes, at partonic level, violate the Froissart bound in the ultraviolet limit. As a matter of fact, although the Wess-Zumino counterterms are introduced in the Lagrangian as dimension-5 local operators to ensure the BRST invariance of the effective action, their contributions to the amplitudes are not sufficient to cancel the divergent behavior of the anomalous poles which affect the BIM amplitude shown in Fig. 16(a). In the supersymmetric generalization of the model that we have presented, this issue of unitarity remains basically the same as for the nonsupersymmetric case.

As we have discussed above, in the latter case the physical axion appears as a massive degree of freedom in the CP -odd sector, due to the presence of a Peccei-Quinn breaking term in the scalar potential. After EWSB the Stückelberg axion b is rotated directly on the physical axion χ and on the two Goldstones $G_Z, G_{Z'}$. Therefore, if we choose the unitary gauge, the only diagram that we can draw in order to erase the bad high energy behavior of


FIG. 15 (color online). The same as Fig. 12 but as a function of g_B , $\tan\beta$ and v_S .

Fig. 16(a) is the second graph (b), where the same amplitude of (a) is mediated by the exchange of the massive axi-Higgs, χ . One can show by a direct study of these two graphs that there is no cancellation of these two contributions at high energy [36]. The problem remains also in the

case of the USSM-A model discussed here. We have again a unitarity bound in the supersymmetric case since the only difference with respect to the nonsupersymmetric case is the contribution of extra fermions circulating in the loops of the BIM amplitude, in particular, the charginos.




FIG. 16. BIM amplitude for $gg \rightarrow \gamma\gamma$ plus the amplitude obtained by the exchange of χ .

XVI. CONCLUSIONS AND PERSPECTIVES

We have presented a generalization of the USSM in the presence of an anomalous $U(1)$ and of a physical axion in the CP -odd scalar sector of the theory, model that we call the USSM-A. This model, which is a direct generalization of a similar construction based on the potential of the MSSM [10], allows Higgs-axion mixing. Both constructions are extensions of a nonsupersymmetric formulation, studied previously [8] (the MLSOM) developed in the context of orientifold vacua of string theory. In the case of the MLSOM, Higgs-axion mixing has been obtained by requiring that the anomalous gauge boson becomes massive by a combination of the Higgs and of the Stückelberg mechanisms, with an axion that is part of the scalar potential. Moving to the supersymmetric case, the generalization of this construction—obtained by using the MSSM superpotential with an extra anomalous $U(1)$ —is found to be characterized by an axino in the spectrum, which appears as a component of the neutralino sector, but not by an axion, since the Stückelberg field does not acquire an axionlike coupling and remains a Goldstone mode. The failure of the MSSM superpotential to provide such a mixing has to be attributed to the structure of the scalar potential of the model. Supersymmetry prohibits a term with a direct presence of the axion in the scalar potential, which otherwise would allow such a mixing.

In our model the mixing occurs indirectly, but the CP -odd sector has to be nonminimal, with an extra singlet which is charged under the anomalous $U(1)$. This approach, as we have emphasized, is quite generic, since its essential working requirement, respect to the MSSM, is the enlargement of the CP -odd sector with one extra SM singlet. Given these minimal requirements, which can be easily satisfied in rather different string vacua, these low energy effective theories capture the essential physical implications of several high energy scenarios, either with a low scale string scale or a much higher scale, as in the heterotic case. Explicit formulations of superpotentials, such as those, for instance, derived from free fermionic models [42], offer the natural ground where to apply the methodology discussed in this work.

Anomalous $U(1)$'s are quite common in string theory but can also be generated, in the corresponding effective Lagrangian, by the decoupling of heavy fermions (and gauge bosons) in grand unified scenarios [41]. It is then

natural to ask what is left at low energy if such decoupling has indeed occurred at some higher scale and it reasonable to foresee that the axion is likely to play a fundamental role [41] in formulating the answer to this question. Clearly, there are corrections to the action discussed in this work, which should be characterized by higher derivative contributions (of dimension larger than 5), i.e. beyond the typical Wess-Zumino terms. Arguments in favor of a possible generalization in this direction of the construction presented in this work have been discussed in previous works [43] and especially in [41]; they are motivated by the fact that anomalies cannot be canceled with local counterterms.

A related issue concerns the size of the mass of the extra Z' in the various models. It is clear that if its decoupling occurs at the Planck scale, then the Stückelberg mass term takes approximately the value of that decoupling scale. This implies that the axionlike couplings induced at low energy are also heavily suppressed. Other interactions, however, in the nonsupersymmetric case, have been found to remain sizeable [41].

A final comment concerns supersymmetry breaking, which may induce phase-dependent terms in the potential. As discussed in [8] for the MLSOM, the axion, in that specific case, gets a sizeable mass which can be as large as the electroweak scale. Similar considerations could remain true in the supersymmetric model that we have presented, although here we have analyzed—by a deliberate choice—the case of a light axion, since we consider this scenario more interesting phenomenologically. In the presence of these phases the pseudoscalar, however, becomes massive. For instance, a mass region of few GeV's is certainly not excluded, as well as a scenario characterized by a very light axion ($\sim 10^{-4}$ eV), and both can be easily included within our analysis. In particular, for an axion in the GeV mass range, for instance, the interactions of this particle are rather similar to those of a light CP -odd Higgs boson, but now with extra interaction with the gauge fields, due to the anomaly, which are not allowed for the rest of the CP -odd sector.

ACKNOWLEDGMENTS

We thank Nikos Irges, Antonio Racioppi and Elisa Manno for discussions. The work of C. C. was supported in part by the European Union through the Marie Curie Research and Training Network Universenet (MRTN-CT-2006-035863) and by The Interreg II Crete-Cyprus program.

APPENDIX A: NOTATIONS

In this appendix we specify our notations. The covariant derivatives are given by

$$\bar{D}_{\dot{A}} = -\bar{\partial}_{\dot{A}} - i\theta^B \sigma_{B\dot{A}}^{\mu} \partial_{\mu} \quad D_A = \partial_A + i\sigma_{AB}^{\mu} \bar{\theta}^{\dot{B}} \partial_{\mu}. \quad (\text{A1})$$

The left/right chiral superfields in terms of field components are given in a generic form as follows

$$\begin{aligned}\hat{\Phi}_L(x, \theta, \bar{\theta}) &= A(x) + i\theta\sigma^\mu\bar{\theta}\partial_\mu A(x) - \frac{1}{4}\theta\theta\bar{\theta}\bar{\theta}\square A(x) \\ &+ \sqrt{2}\theta\psi(x) + \frac{i}{\sqrt{2}}\theta\theta\sigma^\mu\bar{\theta}\partial_\mu\psi(x) \\ &+ \theta\theta F(x),\end{aligned}\quad (\text{A2})$$

$$\begin{aligned}\hat{\Phi}_R^\dagger(x, \theta, \bar{\theta}) &= A^*(x) - i\theta\sigma^\mu\bar{\theta}\partial_\mu A^*(x) - \frac{1}{4}\theta\theta\bar{\theta}\bar{\theta}\square A^*(x) \\ &+ \sqrt{2}\bar{\theta}\bar{\psi}(x) - \frac{i}{\sqrt{2}}\bar{\theta}\bar{\theta}\theta\sigma^\mu\partial_\mu\bar{\psi}(x) \\ &+ \bar{\theta}\bar{\theta}F^*(x).\end{aligned}\quad (\text{A3})$$

A generic scalar superfield \hat{V} in the Wess-Zumino gauge is given by

$$\begin{aligned}\hat{V}(x, \theta, \bar{\theta}) &= \theta\sigma^\mu\bar{\theta}[V_\mu(x) - \partial_\mu B(x)] + \theta\theta\bar{\theta}\bar{\lambda}(x) \\ &+ \bar{\theta}\bar{\theta}\theta\lambda(x) + \theta\theta\bar{\theta}\bar{\theta}d(x)\end{aligned}\quad (\text{A4})$$

where $B(x)$ is a generic real-valued scalar field. The generic expressions for the field-strengths are

$$\begin{aligned}W_\alpha^Y &= -\frac{1}{4}\bar{D}\bar{D}D_\alpha\hat{Y}, & W_\alpha^B &= -\frac{1}{4}\bar{D}\bar{D}D_\alpha\hat{B}, \\ W_\alpha &= -\frac{1}{8g_2}\bar{D}\bar{D}e^{-2g_2\hat{W}}D_\alpha e^{2g_2\hat{W}}, \\ \mathcal{G}_\alpha &= -\frac{1}{8g_s}\bar{D}\bar{D}e^{-2g_s\hat{G}}D_\alpha e^{2g_s\hat{G}}\end{aligned}\quad (\text{A5})$$

where we have used $\hat{W} = \tau^i\hat{W}^i$ with τ^i being the $SU(2)$ generators, while $\hat{G} = T^a\hat{G}^a$ with T^a being the $SU(3)$ generators. The nonsupersymmetric field strength are defined as

$$\begin{aligned}F_{\mu\nu}^Y &= \partial_\mu A_\nu^Y - \partial_\nu A_\mu^Y, & F_{\mu\nu}^B &= \partial_\mu B_\nu - \partial_\nu B_\mu, \\ W_{\mu\nu}^i &= \partial_\mu W_\nu^i - \partial_\nu W_\mu^i - g_2\epsilon^{ijk}W_\mu^j W_\nu^k \\ \mathcal{G}_{\mu\nu}^a &= \partial_\mu G_\nu^a - \partial_\nu G_\mu^a - g_s f^{abc}G_\mu^b G_\nu^c\end{aligned}\quad (\text{A6})$$

APPENDIX B: THE USSM LAGRANGIAN

For completeness we introduce in what follows the USSM Lagrangian that is a part of the total Lagrangian given by $\mathcal{L}_{\text{Tot}} = \mathcal{L}_{\text{USSM}} + \mathcal{L}_{\text{axion}} + \mathcal{L}_{\text{CS}}$.

$$\begin{aligned}\mathcal{L}_{\text{USSM}} &= \mathcal{L}_{\text{lep}} + \mathcal{L}_{\text{quark}} + \mathcal{L}_{\text{Higgs}} + \mathcal{L}_{\text{gauge}} \\ &+ \mathcal{L}_{\text{SMT}} + \mathcal{L}_{\text{GMT}}\end{aligned}\quad (\text{B1})$$

TABLE III. Superfields and their components.

Superfield	Bosonic	Fermionic	Auxiliary
$\hat{\mathbf{b}}(x, \theta, \bar{\theta})$	$b(x)$	$\psi_{\mathbf{b}}(x)$	$F_{\mathbf{b}}(x)$
$\hat{S}(x, \theta, \bar{\theta})$	$S(x)$	$\hat{S}(x)$	$F_S(x)$
$\hat{L}(x, \theta, \bar{\theta})$	$\tilde{L}(x)$	$L(x)$	$F_L(x)$
$\hat{R}(x, \theta, \bar{\theta})$	$\tilde{R}(x)$	$\tilde{R}(x)$	$F_R(x)$
$\hat{Q}(x, \theta, \bar{\theta})$	$\tilde{Q}(x)$	$Q(x)$	$F_Q(x)$
$\hat{U}_R(x, \theta, \bar{\theta})$	$\tilde{U}_R(x)$	$\tilde{U}_R(x)$	$F_{U_R}(x)$
$\hat{D}_R(x, \theta, \bar{\theta})$	$\tilde{D}_R(x)$	$\tilde{D}_R(x)$	$F_{D_R}(x)$
$\hat{H}_1(x, \theta, \bar{\theta})$	$H_1(x)$	$\hat{H}_1(x)$	$F_{H_1}(x)$
$\hat{H}_2(x, \theta, \bar{\theta})$	$H_2(x)$	$\hat{H}_2(x)$	$F_{H_2}(x)$
$\hat{B}(x, \theta, \bar{\theta})$	$B_\mu(x)$	$\lambda_B(x), \bar{\lambda}_B(x)$	$D_B(x)$
$\hat{Y}(x, \theta, \bar{\theta})$	$A_Y^Y(x)$	$\lambda_Y(x), \bar{\lambda}_Y(x)$	$D_Y(x)$
$\hat{W}^i(x, \theta, \bar{\theta})$	$W_\mu^i(x)$	$\lambda_{W^i}(x), \bar{\lambda}_{W^i}(x)$	$D_{W^i}(x)$
$\hat{G}^a(x, \theta, \bar{\theta})$	$G_\mu^a(x)$	$\lambda_{g^a}(x), \bar{\lambda}_{g^a}(x)$	$D_{G^a}(x)$

$$\mathcal{L}_{\text{lep}} = \int d^4\theta [\hat{L}^\dagger e^{2g_2\hat{W}+g_Y\hat{Y}+g_B\hat{B}}\hat{L} + \hat{R}^\dagger e^{2g_2\hat{W}+g_Y\hat{Y}+g_B\hat{B}}\hat{R}]\quad (\text{B2})$$

$$\begin{aligned}\mathcal{L}_{\text{quark}} &= \int d^4\theta [\hat{Q}^\dagger e^{2g_s\hat{G}+2g_2\hat{W}+g_Y\hat{Y}+g_B\hat{B}}\hat{Q} \\ &+ \hat{U}_R^\dagger e^{2g_s\hat{G}+g_Y\hat{Y}+g_B\hat{B}}\hat{U}_R + \hat{D}_R^\dagger e^{2g_s\hat{G}+g_Y\hat{Y}+g_B\hat{B}}\hat{D}_R]\end{aligned}\quad (\text{B3})$$

$$\begin{aligned}\mathcal{L}_{\text{Higgs}} &= \int d^4\theta [\hat{H}_1^\dagger e^{2g_2\hat{W}+g_Y\hat{Y}+g_B\hat{B}}\hat{H}_1 \\ &+ \hat{H}_2^\dagger e^{2g_2\hat{W}+g_Y\hat{Y}+g_B\hat{B}}\hat{H}_2 + \hat{S}^\dagger e^{g_B\hat{B}}\hat{S} \\ &+ \mathcal{W}\delta^2(\bar{\theta}) + \bar{\mathcal{W}}\delta^2(\theta)]\end{aligned}\quad (\text{B4})$$

$$\begin{aligned}\mathcal{L}_{\text{gauge}} &= \frac{1}{4}\int d^4\theta [\mathcal{G}^\alpha\mathcal{G}_\alpha + W^\alpha W_\alpha + W^{Y\alpha}W_\alpha^Y \\ &+ W^{B\alpha}W_\alpha^B]\delta^2(\bar{\theta}) + \text{H.c.}\end{aligned}\quad (\text{B5})$$

$$\begin{aligned}\mathcal{L}_{\text{SMT}} &= -\int d^4\theta\delta^4(\theta, \bar{\theta}) [M_L^2\hat{L}^\dagger\hat{L} + m_R^2\hat{R}^\dagger\hat{R} + M_Q^2\hat{Q}^\dagger\hat{Q} \\ &+ m_U^2\hat{U}_R^\dagger\hat{U}_R + m_D^2\hat{D}_R^\dagger\hat{D}_R + m_1^2\hat{H}_1^\dagger\hat{H}_1 + m_2^2\hat{H}_2^\dagger\hat{H}_2 \\ &+ m_S^2\hat{S}^\dagger\hat{S} + (a_\lambda\hat{S}\hat{H}_1 \cdot \hat{H}_2 + \text{H.c.}) \\ &+ (a_e\hat{H}_1 \cdot \hat{L}\hat{R} + \text{H.c.}) + (a_d\hat{H}_1 \cdot \hat{Q}\hat{D}_R \\ &+ \text{H.c.}) + (a_u\hat{H}_2 \cdot \hat{Q}\hat{U}_R + \text{H.c.})]\end{aligned}\quad (\text{B6})$$

$$\begin{aligned}\mathcal{L}_{\text{GMT}} &= \int d^4\theta \left[\frac{1}{2}(M_G\mathcal{G}^\alpha\mathcal{G}_\alpha + M_W W^\alpha W_\alpha \right. \\ &\left. + M_Y W^{Y\alpha} W_\alpha^Y + M_B W^{B\alpha} W_\alpha^B) + \text{H.c.} \right] \delta^4(\theta, \bar{\theta})\end{aligned}\quad (\text{B7})$$

APPENDIX C: THE O^X MATRIX

$$\begin{aligned}
 O_{11}^X &= \frac{v_2 v_S}{\sqrt{v_1^2 v_2^2 + v_S^2 v^2}}, & O_{12}^X &= \frac{v_1 v_S}{\sqrt{v_1^2 v_2^2 + v_S^2 v^2}}, & O_{13}^X &= \frac{v_1 v_2}{\sqrt{v_1^2 v_2^2 + v_S^2 v^2}}, & O_{14}^X &= 0, \\
 O_{21}^X &= -\frac{v_1(f_1 - 2B_{H_1} g_B x_B + \sqrt{f_1^2 + 4g^2 x_B^2})}{2x_B} \sqrt{\frac{1}{8} - \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, \\
 O_{22}^X &= \frac{v_2(f_1 + 2B_{H_2} g_B x_B + \sqrt{f_1^2 + 4g^2 x_B^2})}{2x_B} \sqrt{\frac{1}{8} - \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, & O_{23}^X &= B_S g_B v_S \sqrt{\frac{1}{8} - \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, \\
 O_{24}^X &= 2M_{st} \sqrt{\frac{1}{8} - \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, & O_{31}^X &= -\frac{v_1(f_1 - 2B_{H_1} g_B x_B - \sqrt{f_1^2 + 4g^2 x_B^2})}{2x_B} \sqrt{\frac{1}{8} + \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, \\
 O_{32}^X &= \frac{v_2(f_1 + 2B_{H_2} g_B x_B - \sqrt{f_1^2 + 4g^2 x_B^2})}{2x_B} \sqrt{\frac{1}{8} + \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, & O_{33}^X &= B_S g_B v_S \sqrt{\frac{1}{8} + \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, \\
 O_{34}^X &= 2M_{st} \sqrt{\frac{1}{8} + \frac{f_1}{8\sqrt{f_1^2 + 4g^2 x_B^2}}}, & O_{41}^X &= \frac{2M_{st} v_1 v_2^2}{\sqrt{(v_1^2 v_2^2 + v^2 v_S^2)[B_S^2 g_B^2 (v_1^2 v_2^2 + v^2 v_S^2) + 4M_{st}^2 v^2]}}, \\
 O_{42}^X &= \frac{2M_{st} v_2 v_1^2}{\sqrt{(v_1^2 v_2^2 + v^2 v_S^2)[B_S^2 g_B^2 (v_1^2 v_2^2 + v^2 v_S^2) + 4M_{st}^2 v^2]}}, \\
 O_{43}^X &= -\frac{2M_{st} v_S v^2}{\sqrt{(v_1^2 v_2^2 + v^2 v_S^2)[B_S^2 g_B^2 (v_1^2 v_2^2 + v^2 v_S^2) + 4M_{st}^2 v^2]}}, & O_{44}^X &= \frac{B_S g_B \sqrt{v_1^2 v_2^2 + v^2 v_S^2}}{\sqrt{B_S^2 g_B^2 (v_1^2 v_2^2 + v^2 v_S^2) + 4M_{st}^2 v^2}}.
 \end{aligned} \tag{C1}$$

-
- [1] M. Ahlers, H. Gies, J. Jaeckel, J. Redondo, and A. Ringwald, Phys. Rev. D **77**, 095001 (2008).
- [2] M. Ahlers, H. Gies, J. Jaeckel, J. Redondo, and A. Ringwald, Phys. Rev. D **76**, 115005 (2007).
- [3] A. De Angelis, O. Mansutti, and M. Roncadelli, Phys. Lett. B **659**, 847 (2008).
- [4] A. De Angelis, O. Mansutti, M. Persic, and M. Roncadelli, arXiv:0807.4246.
- [5] A. De Angelis, O. Mansutti, and M. Roncadelli, Phys. Rev. D **76**, 121301 (2007).
- [6] Z. Berezhiani, L. Gianfagna, and M. Giannotti, Phys. Lett. B **500**, 286 (2001).
- [7] M. I. Vysotsky, Ya. B. Zeldovich, M. Yu. Khlopov, and V. M. Chechetkin, Pis'ma Zh. Eksp. Teor. Fiz. **27**, 533 (1978) [JETP Lett. **27**, 502 (1978)]; Z. G. Berezhiani, M. Yu. Khlopov, and R. R. Khomeriki, Yad. Fiz. **52**, 104 (1990) [Sov. J. Nucl. Phys. **52**, 65 (1990)]; Z. G. Berezhiani, A. S. Sakharov, and M. Yu. Khlopov, Yad. Fiz. **55**, 1918 (1992) [Sov. J. Nucl. Phys. **55**, 1063 (1992)].
- [8] C. Corianò, N. Irges, and E. Kiritsis, Nucl. Phys. **B746**, 77 (2006).
- [9] E. Kiritsis, Fortschr. Phys. **52**, 200 (2004).
- [10] P. Anastasopoulos, F. Fucito, A. Lionetto, G. Pradisi, A. Racioppi, and Y. S. Stanev, Phys. Rev. D **78**, 085014 (2008).
- [11] J. De Rydt, J. Rosseel, Torsten T. Schmidt, A. Van Proeyen, and M. Zagermann, Classical Quantum Gravity **24**, 5201 (2007).
- [12] M. Zagermann, arXiv:0801.1666.
- [13] R. Armillis, C. Corianò, M. Guzzi, and S. Morelli, Nucl. Phys. **B814**, 15679 (2009).
- [14] B. Kors and P. Nath, J. High Energy Phys. **12** (2004) 005.
- [15] C. Corianò, M. Guzzi, N. Irges, and A. Mariano, Phys. Lett. B **671**, 87 (2009).
- [16] J. E. Kim and H. P. Nilles, Phys. Lett. B **138B**, 150 (1984).
- [17] V. Jain and R. Shrock, Phys. Lett. B **352**, 83 (1995).
- [18] Y. Nir, Phys. Lett. B **354**, 107 (1995).
- [19] D. Suematsu and Y. Yamagishi, Int. J. Mod. Phys. A **10**, 4521 (1995).
- [20] M. Cvetič, Durmus A. Demir, J. R. Espinosa, L. L. Everett,

- and P. Langacker, Phys. Rev. D **56**, 2861 (1997).
- [21] M. Cvetič and P. Langacker, Phys. Rev. D **54**, 3570 (1996).
- [22] C. Balazs, M. S. Carena, A. Freitas, and C. E. M. Wagner, J. High Energy Phys. 06 (2007) 066.
- [23] U. Ellwanger and C. Hugonie, arXiv:hep-ph/0006222.
- [24] P. Anastasopoulos, M. Bianchi, E. Dudas, and E. Kiritsis, J. High Energy Phys. 11 (2006) 057.
- [25] D. Feldman, Z. Liu, and P. Nath, J. High Energy Phys. 11 (2006) 007.
- [26] D. Feldman, Z. Liu, and P. Nath, Phys. Rev. D **75**, 115001 (2007).
- [27] C. Corianò, M. Guzzi, and A. Mariano (unpublished).
- [28] W. Fischler, H. P. Nilles, J. Polchinski, S. Raby, and L. Susskind, Phys. Rev. Lett. **47**, 757 (1981).
- [29] J. Preskill, Ann. Phys. (N.Y.) **210**, 323 (1991).
- [30] C. Corianò, N. Irges, and S. Morelli, J. High Energy Phys. 07 (2007) 008.
- [31] C. Panagiotakopoulos and K. Tamvakis, Phys. Lett. B **446**, 224 (1999).
- [32] S. A. Abel, S. Sarkar, and P. L. White, Nucl. Phys. **B454**, 663 (1995).
- [33] D. J. Miller, R. Nevzorov, and P. M. Zerwas, Nucl. Phys. **B681**, 3 (2004).
- [34] C. Corianò, N. Irges, and S. Morelli, Nucl. Phys. **B789**, 133 (2008).
- [35] R. Armillis, C. Corianò, and M. Guzzi, J. High Energy Phys. 05 (2008) 015.
- [36] C. Corianò, M. Guzzi, and S. Morelli, Eur. Phys. J. C **55**, 629 (2008).
- [37] R. Armillis, C. Corianò, M. Guzzi, and S. Morelli, J. High Energy Phys. 10 (2008) 034.
- [38] J. Kalinowski, S. F. King, and J. P. Roberts, J. High Energy Phys. 01 (2009) 066.
- [39] C. Amsler *et al.*, Phys. Lett. B **667**, 1 (2008).
- [40] A. Lionetto and A. Racioppi, arXiv:0905.4607; F. Fucito, A. Lionetto, A. Mammarella, and A. Racioppi, arXiv:0811.1953.
- [41] C. Corianò and M. Guzzi, arXiv:0905.4462.
- [42] A. E. Faraggi, E. Manno, and C. Timirgaziu, Eur. Phys. J. C **50**, 701 (2007).
- [43] C. Corianò and N. Irges, Phys. Lett. B **651**, 298 (2007).