

## Dirichlet Series and Gamma Function Associated with Rational Functions

by

Mauro SPREAFICO

*Presented by Jerzy KACZOROWSKI*

**Summary.** We investigate zeta regularized products of rational functions. As an application, we obtain the asymptotic expansion of the Euler Gamma function associated with a rational function.

**1. Introduction.** Let  $r(z) = cp_h(z)/q_k(z)$  be a rational function of  $z$ , where  $p_h$  and  $q_k$  are monic polynomials with real coefficients of degree  $h$  and  $k$ , respectively, and  $c \neq 0$  is a real number. Factoring  $r(z)$  into the product

$$r(z) = c \frac{p_h(z)}{q_k(z)} = c \frac{(z + a_1) \cdots (z + a_h)}{(z + b_1) \cdots (z + b_k)},$$

it is clear (see for example [8, 12.13]) that the infinite product

$$\prod_{n=1}^{\infty} c \frac{p_h(n)}{q_k(n)}$$

converges if  $c = 1$ ,  $h = k$ ,  $a_1 + \cdots + a_h - b_1 - \cdots - b_h = 0$ , and assuming that no factor in the denominator vanishes. If this is the case, it is a result of Euler that

$$\prod_{n=1}^{\infty} \frac{p_h(n)}{q_h(n)} = \frac{\Gamma(1 + b_1) \cdots \Gamma(1 + b_h)}{\Gamma(1 + a_1) \cdots \Gamma(1 + a_h)}.$$

M. Eie [4, Main theorem II] proved that this result generalizes to zeta regularized products when  $q_k(z) = 1$  and  $|a_l| < 1$ . Recall that if  $\Lambda = \{\lambda_n\}_{n=1}^{\infty}$  is a sequence of complex numbers with a unique accumulation point at infinity

---

2010 *Mathematics Subject Classification*: Primary 11M06.

*Key words and phrases*: rational function, Riemann zeta function, zeta regularized product.

ity and genus  $g$  (see for example [2, 7.5], or [6, Section 2] for the definition), and if  $\Lambda$  is contained in some suitable sector of the complex plane, then the *zeta regularization* of the infinite product

$$\prod_{n=1}^{\infty} \lambda_n$$

is by definition  $e^{-\zeta'(0, \Lambda)}$ , where the zeta function associated to  $\Lambda$  is defined by the Dirichlet series

$$\zeta(s, \Lambda) = \sum_{n=1}^{\infty} \lambda_n^{-s}$$

when  $\text{Re}(s) > g$ , and by analytic continuation elsewhere, and where by  $\zeta'(0, \Lambda)$  we mean the finite part of  $\zeta(s, \Lambda)$  if  $\zeta(s, \Lambda)$  has a pole at  $s = 0$  (we refer to [6] for details). If the unique accumulation point of  $\Lambda$  is zero, then we define the associated zeta function by  $\zeta(s, \Lambda) = \zeta(-s, 1/\Lambda)$ .

If  $\Lambda = \{cp_h(n)/q_k(n)\}_{n=1}^{\infty}$ , we denote by  $\zeta(s, cp_h/q_k)$  the associated zeta function, and we call it the *zeta function associated with a rational function*. The polynomial zeta function  $\zeta(s, p_k)$  has been studied in the cited work of Eie. Subsequently, the construction has been generalized by studying multiple polynomial zeta functions in [5], and introducing polynomial multiplicity in [3].

In this note, we extend this construction to the case of rational functions. Our first result is the following proposition, which also gives an elementary proof of Main Theorem II in [4].

**PROPOSITION 1.** *Let  $c \neq 0$ , and  $a_1, \dots, a_h$  and  $b_1, \dots, b_k$  be complex numbers with  $|a_j| < 1$ ,  $|b_j| < 1$ , and  $h \neq k$ . Then the zeta regularization of the infinite product*

$$\prod_{n=1}^{\infty} c \frac{p_h(n)}{q_k(n)} = \prod_{n=1}^{\infty} c \frac{(n + a_1) \cdots (n + a_h)}{(n + b_1) \cdots (n + b_k)}$$

is

$$e^{-\zeta'(0, cp_h/q_k)} = (2\pi)^{\frac{h-k}{2}} c^{-\frac{a_1 + \dots + a_h - b_1 - \dots - b_k}{h-k} - \frac{1}{2}} \frac{\Gamma(1 + b_1) \cdots \Gamma(1 + b_k)}{\Gamma(1 + a_1) \cdots \Gamma(1 + a_h)}.$$

As a second result, we present the following natural application of Proposition 1. Define the Euler Gamma function associated with the rational function  $r(z) = cp_h(z)/q_k(z)$ , with  $h > k > 0$ , to be the Weierstrass product

$$\Gamma(z, cp_h/q_k) = \prod_{n=1}^{\infty} \frac{e^{gz/r(n)}}{1 + z/r(n)},$$

where we put  $g = 1$  if  $h = k + 1$ , and  $g = 0$  otherwise (note that  $g$  is the genus of the sequence  $\Lambda$ ). Then we have the following asymptotic expansion,

where the notation  $\text{Res}_{s=s_0} f(s)$  denotes the coefficient of the term  $(s - s_0)^{-l}$  in the Laurent expansion of  $f(s)$  at  $s = s_0$  (see for example [1, p. 420]).

PROPOSITION 2. For large  $z$  with  $|\arg(z)| < \pi$ ,

$$\log \Gamma(z, cp_h/q_k) = \begin{cases} \frac{\pi c^{\frac{1}{k-h}}}{\sin \frac{\pi}{k-h}} z^{\frac{1}{h-k}} & \text{if } h > k + 1, \\ \frac{1}{c} z \log z + \left( \text{Res}_{s=1} \zeta(s, cp_h/q_k) - \frac{1}{c} \right) z & \text{if } h = k + 1, \\ + \left( \frac{1}{2} + \frac{a_1 + \dots + a_h - b_1 - \dots - b_k}{h - k} \right) \log z \\ - \left( \frac{1}{2} + \frac{a_1 + \dots + a_h - b_1 - \dots - b_k}{h - k} \right) \log c \\ + \frac{h - k}{2} \log 2\pi + \log \frac{\Gamma(1 + b_1) \cdots \Gamma(1 + b_k)}{\Gamma(1 + a_1) \cdots \Gamma(1 + a_h)} + o(1). \end{cases}$$

The proofs of these propositions are presented in the next two sections.

**2. The proof of Proposition 1.** Expanding the powers of the binomials we obtain, for large  $\text{Re}(s)$ ,

$$c^s \zeta(s, cp_h/q_k) = \sum_{j,l=0}^{\infty} \binom{-s}{j} \binom{s}{l} a^j b^l \zeta((h-k)s + |j| + |l|),$$

where  $\zeta(s)$  is the Riemann zeta function, we use the multi-indices  $j = (j_1, \dots, j_h)$ ,  $l = (l_1, \dots, l_k)$ , and  $|j| = j_1 + \dots + j_h$ ,  $|l| = l_1 + \dots + l_k$ ,  $a_j^j = a_1^{j_1} \cdots a_h^{j_h}$ ,  $b_k^k = b_1^{l_1} \cdots b_k^{l_k}$ ,  $\binom{-s}{j} = \binom{-s}{j_1} \cdots \binom{-s}{j_h}$ ,  $\binom{s}{l} = \binom{s}{l_1} \cdots \binom{s}{l_k}$ . Thus,

$$\begin{aligned} c^s \zeta(s, cp_h/q_k) &= \zeta((h-k)s) + \sum_{\alpha=1}^h \sum_{j_\alpha=1}^{\infty} \binom{-s}{j_\alpha} a_\alpha^{j_\alpha} \zeta((h-k)s + j_\alpha) \\ &\quad + \sum_{\beta=1}^k \sum_{l_\beta=1}^{\infty} \binom{s}{l_\beta} b_\beta^{l_\beta} \zeta((h-k)s + l_\beta) + \varphi(s), \end{aligned}$$

where  $\varphi(s)$  has a zero of degree 2 at  $s = 0$ . Isolating the singular terms, we obtain

$$\begin{aligned} (1) \quad c^s \zeta(s, cp_h/q_k) &= \zeta((h-k)s) - s \zeta((h-k)s + 1) \left( \sum_{\alpha=1}^h a_\alpha - \sum_{\beta=1}^k b_\beta \right) \\ &\quad + \sum_{\alpha=1}^h \sum_{j_\alpha=2}^{\infty} \binom{-s}{j_\alpha} a_\alpha^{j_\alpha} \zeta((h-k)s + j_\alpha) \\ &\quad + \sum_{\beta=1}^k \sum_{l_\beta=2}^{\infty} \binom{s}{l_\beta} b_\beta^{l_\beta} \zeta((h-k)s + l_\beta) + \varphi(s). \end{aligned}$$

The analytic continuation at  $s = 0$  of the function on the right side of (1) is given by that of the Riemann zeta function, and is regular at  $s = 0$ . In particular,

$$\begin{aligned} \frac{d}{ds} \Big|_{s=0} \sum_{j_\alpha=2}^{\infty} \binom{-s}{j_\alpha} \zeta((h-k)s + j_\alpha) a_\alpha^{j_\alpha} &= \sum_{j_\alpha=2}^{\infty} \frac{(-1)^{j_\alpha}}{j_\alpha} \zeta(j_\alpha) a_\alpha^{j_\alpha} \\ &= \log \Gamma(1 + a_\alpha) + \gamma a_\alpha, \\ \frac{d}{ds} \Big|_{s=0} \sum_{l_\beta=2}^{\infty} \binom{s}{l_\beta} \zeta((h-k)s + l_\beta) b_\beta^{l_\beta} &= - \sum_{l_\beta=2}^{\infty} \frac{(-1)^{l_\beta}}{l_\beta} \zeta(l_\beta) b_\beta^{l_\beta} \\ &= - \log \Gamma(1 + b_\beta) - \gamma b_\beta. \end{aligned}$$

This gives the expansions near  $s = 0$  of the different terms:

$$\begin{aligned} -s \zeta((h-k)s + 1) \left( \sum_{\alpha=1}^h a_\alpha - \sum_{\beta=1}^k b_\beta \right) &= - \frac{1}{h-k} \left( \sum_{\alpha=1}^h a_\alpha - \sum_{\beta=1}^k b_\beta \right) \\ &\quad - \gamma \left( \sum_{\alpha=1}^h a_\alpha - \sum_{\beta=1}^k b_\beta \right) s + O(s^2), \\ \sum_{\alpha=1}^h \sum_{j_\alpha=2}^{\infty} \binom{-s}{j_\alpha} a_\alpha^{j_\alpha} \zeta((h-k)s + j_\alpha) &= \left( \log \prod_{\alpha=1}^h \Gamma(1 + a_\alpha) + \gamma \sum_{\alpha=1}^h a_\alpha \right) s + O(s^2), \\ \sum_{\beta=1}^k \sum_{l_\beta=2}^{\infty} \binom{s}{l_\beta} b_\beta^{l_\beta} \zeta((h-k)s + l_\beta) &= - \left( \log \prod_{\beta=1}^k \Gamma(1 + b_\beta) + \gamma \sum_{\beta=1}^k b_\beta \right) s + O(s^2), \end{aligned}$$

and hence

$$\begin{aligned} \zeta(0, p_h/q_k) &= \zeta(0) - \frac{1}{h-k} \left( \sum_{\alpha=1}^h a_\alpha - \sum_{\beta=1}^k b_\beta \right), \\ \zeta(0, p_h/q_k) &= (h-k) \zeta'(0) + \log \frac{\prod_{\alpha=1}^h \Gamma(1 + a_\alpha)}{\prod_{\beta=1}^k \Gamma(1 + b_\beta)}. \end{aligned}$$

**3. The proof of Proposition 2.** We shall use some notations and results from [6] and [7], concerning sequences of spectral type. Let us indicate the main steps of the proof.

Let  $S$  be any sequence of positive real numbers, which is simple regular of spectral type with genus  $g$ . Let  $F(z, S)$  denote the Fredholm determinant associated to  $S$  (see [6, p. 866]; also note that its inverse is called the Gamma function in [7]). There exists an expansion (use Definition 2.1 and Lemma 2.7 in [6] or Definitions 2.1 and 2.7 in [7])

$$-\log F(z, S) = \sum_{j=0}^{\mathbf{g}} a_{j,1} z^j \log z + \sum_{j=0}^J a_{\alpha_j,0} z^{\alpha_j} + o(z^{\alpha_J})$$

for large  $z$  with  $|\arg(z)| < \pi$  and  $\alpha_0 > \alpha_1 > \dots > \alpha_J$ . Observe that our sequence  $\Lambda$  is a simply regular sequence of spectral type with genus  $\mathbf{g}$ , where  $\mathbf{g} = 1$  if  $h = k + 1$  and  $\mathbf{g} = 0$  otherwise. Indeed,  $\Lambda$  is a sequence of spectral type by Lemma 2.5 in [6], and is simply regular by Proposition 2.11 in [7], since the possible poles of the zeta function  $\zeta(s, \Lambda)$  are at most simple. Moreover, by the same proposition the possible poles of the zeta function are located at  $s = \alpha_j$ , and by Remark 2.9 in [7],  $\alpha_0 < \mathbf{g} + 1$ . Now, using the expansion given in the previous section, it is easy to see that  $\zeta(s, \Lambda)$  has at most one simple pole on the positive part of the real axis, and this pole is at  $s = 1$  if  $\mathbf{g} = 1$ , and at  $s = 1/(h - k)$  otherwise. It follows that the unique possible positive value of the  $\alpha_j$  is either  $\alpha_0 = 1$ , if  $\mathbf{g} = 1$ , or  $\alpha_0 = \frac{1}{h-k}$ , if  $\mathbf{g} = 0$ ; and that  $\alpha_1 = 0$  for any  $\mathbf{g}$ . Also note that  $\log F(z, \Lambda) = -\log \Gamma(z, \Lambda)$ . This means that

$$\log \Gamma(z, \Lambda) = \sum_{j=0}^{\mathbf{g}} a_{j,1} z^j \log z + \sum_{j=0}^1 a_{\alpha_j,0} z^{\alpha_j} + o(1).$$

The values of  $a_{x,k}$ 's can be calculated explicitly as follows (see Propositions 2.11 and 2.14 in [7], and Proposition 2.6 in [6]):

$$\begin{aligned} a_{0,0} &= -\operatorname{Res}_0 \zeta'(s, \Lambda), \\ a_{0,1} &= -\operatorname{Res}_0 \zeta(s, \Lambda), \\ a_{\alpha_0,0} &= \begin{cases} a_{\frac{1}{h-k},0} = \Gamma\left(\frac{1}{h-k}\right) \Gamma\left(\frac{1}{k-h}\right) \operatorname{Res}_{s=\frac{1}{h-k}} \zeta(s, \Lambda) = \frac{\pi c^{\frac{1}{k-h}}}{\sin \frac{\pi}{k-h}}, & \mathbf{g} = 0, \\ a_{1,0} = \operatorname{Res}_0 \zeta(s, \Lambda) - \operatorname{Res}_1 \zeta(s, \Lambda) = \operatorname{Res}_0 \zeta(s, \Lambda) - \frac{1}{c}, & \mathbf{g} = 1, \end{cases} \\ a_{\alpha_0,1} &= \begin{cases} a_{\frac{1}{h-k},1} = 0, & \mathbf{g} = 0, \\ a_{1,1} = \operatorname{Res}_1 \zeta(s, \Lambda) = \frac{1}{c}, & \mathbf{g} = 1. \end{cases} \end{aligned}$$

Applying Proposition 1, we are done.

**4. Remarks.** In this section we investigate the case  $h = k$ . Before, we discuss multiplicativity of zeta regularization. This appears in the interpretation of the infinite product  $\prod \lambda_n$  as the determinant of the infinite diagonal matrix  $\Lambda$  with entries  $\lambda_n$ . Namely, we set

$$\det_{\zeta} \Lambda = e^{-\zeta'(0, \Lambda)}.$$

Then it is natural to ask: given two infinite diagonal matrices  $A_1$  and  $A_2$ , is  $\det_{\zeta} A_1 A_2 = \det_{\zeta} A_1 \det_{\zeta} A_2$ ? Multiplicativity of determinants clearly corresponds to additivity of the derivative at zero of the associated zeta functions. Now, it is clear that  $\zeta(0, cA) = \zeta(0, A)$  for any  $c \neq 0$ , and that

$$\zeta'(0, cA) = -\zeta(0, A) \log c + \zeta'(0, A).$$

Restricting to the case where the  $\lambda_n$  are rational functions of  $n$ , it follows from Proposition 1 and the formula

$$\zeta(0, p_h) = -\frac{1}{2} - \frac{1}{h}(a_1 + \cdots + a_h)$$

that  $\zeta'(0, c_1 c_2 p_{1, h_1} p_{2, h_2})$  is not equal to  $\zeta'(0, c_1 p_{1, h_1}) + \zeta'(0, c_2 p_{2, h_2})$  (however, note this is the case when  $h_1 = h_2$  and  $p_{1, h_1} = p_{2, h_2}$ ). Therefore, we further restrict to monic polynomials, and in this case it is easy to see that (if  $h_j \neq k_j$ )

$$\zeta'(0, p_{1, h_1} p_{2, h_2} / q_{1, k_1} q_{2, k_2}) = \zeta'(0, p_{1, h_1} / q_{1, k_1}) + \zeta'(0, p_{2, h_2} / q_{2, k_2}).$$

Thus we consider the case  $k = h$  only for monic polynomials. It is clear that a zeta function for the sequence  $\{p_h(n)/q_h(n)\}_{n=1}^{\infty}$  cannot be defined, since the exponent of convergence is not finite. However, we can introduce the following regularization of the infinite product: we define the regularized product

$$\begin{aligned} \prod_{n=1}^{\infty} \frac{p_h(n)}{q_h(n)} &= \prod_{n=1}^{\infty} \frac{(n+a_1) \cdots (n+a_h)}{(n+b_1) \cdots (n+b_h)} \\ &= e^{a_1 + \cdots + a_h - b_1 - \cdots - b_h} \prod_{n=1}^{\infty} \frac{(n+a_1) \cdots (n+a_h)}{(n+b_1) \cdots (n+b_h)} e^{-(a_1 + \cdots + a_h - b_1 - \cdots - b_h)/n}. \end{aligned}$$

It is then easy to see that the product on the right side converges, as desired, to

$$\frac{\Gamma(1+b_1) \cdots \Gamma(1+b_h)}{\Gamma(1+a_1) \cdots \Gamma(1+a_h)} = \frac{e^{-\zeta'(0, p_h)}}{e^{-\zeta'(0, q_h)}}.$$

We conclude by observing that the method described in this note can be used to obtain a formula for the derivative at zero of the zeta function studied by Dąbrowski in [3], where some polynomial multiplicity has been introduced.

**Acknowledgments.** We thank the referee for his comments and remarks to strengthen the presentation of the results in this note.

#### References

- [1] J. Brüning and R. Seeley, *The resolvent expansion for second order regular singular operators*, J. Funct. Anal. 73 (1988), 369–415.

- 
- [2] E. T. Copson, *Theory of Functions of a Complex Variable*, Oxford Univ. Press, 1935.
  - [3] A. Dąbrowski, *On zeta functions associated with polynomials*, Bull. Austral. Math. Soc. 61 (2000), 455–458.
  - [4] M. Eie, *On Dirichlet series associated with a polynomial*, Proc. Amer. Math. Soc. 110 (1990), 583–590.
  - [5] M. Eie and K.-W. Chen, *A theorem on zeta functions associated with polynomials*, Trans. Amer. Math. Soc. 351 (1999), 3217–3228.
  - [6] M. Spreafico, *Zeta invariants for sequences of spectral type, special functions and the Lerch formula*, Proc. Roy. Soc. Edinburgh 281 (2006), 865–889.
  - [7] —, *Zeta invariants for double sequences of spectral type*, arXiv:math/0607816.
  - [8] E. T. Whittaker and G. N. Watson, *Modern Analysis*, Cambridge Univ. Press, 1963.

Mauro Spreafico  
ICMC, Universidade de São Paulo  
São Carlos, Brazil  
E-mail: mauros@icmc.usp.br

*Received October 15, 2010;*  
*received in final form November 1, 2010*

(7797)

